

Diversity in Short β -Peptide 12-Helices: High-Resolution Structural Analysis in Aqueous Solution of a Hexamer Containing Sulfonylated Pyrrolidine Residues

Hee-Seung Lee, Faisal A. Syud, Xifang Wang, and Samuel H. Gellman*

Department of Chemistry, University of Wisconsin
Madison, Wisconsin 53706

Received March 20, 2001

Oligomers that adopt well-defined conformations in solution (“foldamers”) have been a subject of increasing interest.¹ Results from many laboratories demonstrate that shape control can be achieved with a wide variety of backbones, and recent efforts have shown that foldamers can be endowed with useful activities.² Many applications require placement of specific functional groups at defined positions along the foldamer backbone, so that folding brings these groups into the arrangement necessary for activity.

Described here is a strategy for functionalizing the 12-helix, a secondary structure defined by 12-membered-ring hydrogen bonds [C=O(*i*) \rightarrow N–H(*i*+3)] that is formed by β -amino acid oligomers in which the residues are constrained by five-membered rings.³ Two appropriately constrained residues have been identified, *trans*-2-aminocyclopentanecarboxylic acid (ACPC)^{3a} and *trans*-3-aminopyrrolidine-4-carboxylic acid (APC),^{3d} each of which can be prepared efficiently in large quantities in either enantiomeric form.⁴ We have now explored sulfonylated APC (S-APC) residues for functionalization of water-soluble 12-helices. Attachment of side chains via N-sulfonylation should be advantageous relative to N-alkylation, which would introduce cationic charge, or N-acylation, which would interfere with structural characterization because of *cis*–*trans* rotamer equilibria of the resulting tertiary amide groups. However, sulfonylation introduces a non-*sp*³ atom⁵ into the five-membered ring of the β -amino acid residue and might therefore adversely affect 12-helix stability by altering the C α –C β torsional preference of S-APC residues relative to APC and ACPC residues.

* To whom correspondence should be addressed (e-mail: gellman@chem.wisc.edu).


(1) Reviews: (a) Seebach, D.; Matthews, J. L. *J. Chem. Soc., Chem. Commun.* **1997**, 2015–2022. (b) Gellman, S. H. *Acc. Chem. Res.* **1998**, *31*, 173. (c) DeGrado, W. F.; Schneider, J. P.; Hamuro, Y. *J. Pept. Res.* **1999**, *54*, 206. (d) Kirshenbaum, K.; Zuckermann, R. N.; Dill, K. A. *Curr. Opin. Struct. Biol.* **1999**, *9*, 530. (e) Stigers, K. D.; Soth, M. J.; Nowick, J. S. *Curr. Opin. Chem. Biol.* **1999**, *3*, 714. (f) Barron, A. E.; Zuckermann, R. N. *Curr. Opin. Chem. Biol.* **1999**, *3*, 681. (g) Gademann, K.; Hintermann, T.; Schreiber, J. V. *Curr. Med. Chem.* **1999**, *6*, 905.

(2) (a) Biologically active β -peptides: Werder, M.; Hausre, H.; Abele, S.; Seebach, D. *Helv. Chim. Acta* **1999**, *82*, 1774. Hamuro, Y.; Schneider, J. P.; DeGrado, W. F. *J. Am. Chem. Soc.* **1999**, *121*, 12200. Porter, E. A.; Wang, X.; Lee, H.-S.; Weisblum, B.; Gellman, S. H. *Nature* **2000**, *404*, 565. (b) A carbohydrate oligomer that binds to DNA: Xuereb, H.; Maletic, M.; Gildersleeve, J.; Pelczar, I.; Kahne, D. *J. Am. Chem. Soc.* **2000**, *122*, 1883. (c) *m*-Phenylene ethynylene oligomers that bind specific guests: Prince, R. B.; Barnes, S. A.; Moore, J. S. *J. Am. Chem. Soc.* **2000**, *122*, 2758. Tanatani, A.; Mio, M. J.; Moore, J. S. *J. Am. Chem. Soc.* **2001**, *123*, 1792.


(3) (a) Appella, D. H.; Christianson, L. A.; Klein, D. A.; Powell, D. R.; Huang, X.; Barchi, J. J.; Gellman, S. H. *Nature* **1997**, *387*, 381. (b) Appella, D. H.; Christianson, L. A.; Klein, D. A.; Richards, M. R.; Powell, D. R.; Gellman, S. H. *J. Am. Chem. Soc.* **1999**, *121*, 7574. (c) Barchi, J. J.; Huang, X.; Appella, D. H.; Christianson, L. A.; Durell, S. R.; Gellman, S. H. *J. Am. Chem. Soc.* **2000**, *122*, 2711. (d) Wang, X.; Espinosa, J. F.; Gellman, S. H. *J. Am. Chem. Soc.* **2000**, *122*, 4821. (e) Christianson, L. A.; Lucero, M. J.; Appella, D. H.; Klein, D. A.; Gellman, S. H. *J. Comput. Chem.* **2000**, *21*, 763. (f) Alternative residues for the 12-helix: Winkler, J. D.; Piatnitski, E. L.; Mehlmann, J.; Kasperec, J.; Axelsen, P. H. *Angew. Chem., Int. Ed.* **2001**, *40*, 743.

(4) (a) Lee, H.-S.; LePlae, P. L.; Porter, E. A.; Gellman, S. H. *J. Org. Chem.* **2001**, *66*, 3597. (b) LePlae, P. L.; Umezawa, N.; Lee, H.-S.; Gellman, S. H. *J. Org. Chem.*, in press.

(5) Radkiewicz, J. L.; McAllister, M. A.; Goldstein, E.; Houk, K. N. *J. Org. Chem.* **1998**, *63*, 1419 and references therein.



Fmoc-protected S-APC residues were prepared via straightforward methods⁶ and used for solid-phase synthesis of hexamers **1**–**3**. The two S-APC residues in **1**–**3** illustrate that alkyl and aromatic side chains can be introduced in this way; a variety of polar S-APC residues should also be readily available (for example, we have prepared a taurine-derived S-APC residue, with which it should be possible to replace the cationic APC residues of **1**–**3**). Hexamer **1** was analyzed by 2D NMR in 9:1 H₂O:D₂O to determine whether a backbone comprised solely of APC and S-APC residues adopts the 12-helical conformation. ¹H chemical shifts of **1** did not change significantly upon dilution from 2.5 mM to 0.3 mM, which suggests that the hexamer does not self-associate in this concentration range. The ¹H NMR resonances of **1** in water are more dispersed than those of previously examined^{3d} APC/ACPC hexamer **4**, presumably because of the greater residue diversity of **1** relative to **4** and ring current effects from the aromatic side chain. Seven NOEs (present in both



NOESY^{7a} and ROESY^{7b} spectra) between protons on nonadjacent residues were observed along the backbone of **1** (Figure 1a). All three types of nonadjacent NOEs, C β H_{*i*} \rightarrow NH_{*i*+2}, C β H_{*i*} \rightarrow C α H_{*i*+2}, and C β H_{*i*} \rightarrow NH_{*i*+3}, are characteristic of the 12-helical folding pattern.^{3a,d} The lack of NOEs in **1** between residues 1 and 3 (numbered from N-terminus) suggests that the N-terminus of the 12-helix is frayed; terminal fraying is observed among α -helices formed by conventional peptides in aqueous solution.⁸

Comparison of NOE data for hexamers **1** and **4** suggests that incorporating S-APC residues (in **1**) in place of ACPC residues (in **4**) leads to subtle differences in 12-helix geometry. Previous data^{3d} showed that **4** in water displayed a set of four weak C β H_{*i*} \rightarrow C α H_{*i*+1} NOEs (only C β H₅ \rightarrow C α H₆ was missing). In contrast, no C β H_{*i*} \rightarrow C α H_{*i*+1} NOEs were observed for **1**, which suggests that the 12-helix formed by **1** is slightly more tightly wound than the 12-helix formed by **4** (Figure 1b). The subtle differences between the 12-helical conformations of **1** and **4** are comparable to differences among α -helices formed by conventional peptides.⁹

NOE data obtained for **1** in 9:1 H₂O:D₂O were used for NOE-restrained dynamics simulations with the program DYANA.¹⁰ This approach was used to generate 400 structures, the best 10 of which (all 12-helical) were used as starting points for NOE-restrained

(6) Please see the Supporting Information.

(7) (a) Macura, S.; Ernst, R. R. *Mol. Phys.* **1980**, *41*, 95. (b) Bothner-By, A. A.; Stephens, R. L.; Lee, J.; Warren, C. D.; Jeanloz, R. W. *J. Am. Chem. Soc.* **1984**, *106*, 811.

(8) Rohl, C. A.; Baldwin, R. L. *Biochemistry* **1994**, *31*, 7760. Scholtz, J. M.; Baldwin, R. L. *Annu. Rev. Biophys. Biomol. Struct.* **1992**, *21*, 95.

(9) For example, ideal α -helices have 3.6 residues per turn while α -helices that participate in coiled-coil structures are slightly unwound, with 3.5 residues per turn: Creighton, T. E. *Proteins: Structures and Molecular Properties*, 2nd ed.; W. H. Freeman and Company: New York, 1993.

(10) Guntert, P.; Mumenthaler, C.; Wüthrich, K. *J. Mol. Biol.* **1997**, *273*, 283. The structure library in DYANA was modified to include cyclopentane and pyrrolidine rings for these calculations.




Figure 1. (a) Graphical summary of NOEs between nonadjacent residues observed for hexa- β -peptide **1** (2.5 mM, trifluoroacetate salt) in 9:1 $\text{H}_2\text{O}:\text{D}_2\text{O}$, 4 °C. Three types of NOE were observed: $\text{C}_\beta\text{H}_i \rightarrow \text{NH}_{i+2}$, $\text{C}_\beta\text{H}_i \rightarrow \text{C}_\alpha\text{H}_{i+2}$, and $\text{C}_\beta\text{H}_i \rightarrow \text{NH}_{i+3}$ (solid arrows in part b). All of these NOEs are consistent with the 12-helical conformation; no NOEs inconsistent with the 12-helical conformation were observed. (b) Cartoon showing the spatial relationship between protons on residue i and protons on residues $i+2$ and $i+3$ in the 12-helix; as mentioned in the text, $\text{C}_\beta\text{H}_i \rightarrow \text{C}_\alpha\text{H}_{i+1}$ NOEs (dashed arrow) were observed for **4** (ref 3d) but not for **1**, which suggests that the helix formed by **1** is slightly more tightly wound.




Figure 2. NOE-restrained dynamics results for hexa- β -peptide **1**, based on NMR data obtained in 9:1 $\text{H}_2\text{O}:\text{D}_2\text{O}$, 4 °C (details in text). Overlay of 10 structures, backbone atoms only, with low energy and NOE restraint violation. The N-termini are at the top.

simulated annealing with the SYBYL 6.4 program (Tripos Software, St. Louis, MO). Figure 2 shows an overlay of the resulting 10 structures, backbone atoms only. Fraying, apparent at both ends, is pronounced at the N-terminus, but the 12-helix is well-formed in the central portion. Among the 10 structures shown root-mean-squared deviation is 0.69 ± 0.51 Å for residues 3–6 (backbone atoms). The 12-helical conformation deduced for **1** in water agrees well with the conformation previously deduced for **4** in methanol^{3d} and with crystal structures of ACPC oligo-

mers.^{3a,b} It was not possible to determine a structure of **4** in water because of overlap among crucial proton resonances and a general decrease in the number of characteristic NOEs, relative to methanol.^{3d} In contrast, the proton resonance dispersion of **1** was comparable in water and methanol, and the number and intensities of the helix-defining NOEs were comparable in the two solvents.

Circular dichroism is widely used to assess the secondary structures of conventional peptides,¹¹ and CD has also been useful for β -peptides^{12,13} and other unnatural oligomers.¹⁴ CD spectra of **1–3** in H_2O and in CH_3OH show a characteristic maximum near 200 nm and a minimum near 220 nm,⁶ which is consistent with theoretical predictions¹² and previously reported CD data³ for 12-helical β -peptides. In CH_3OH the maximum and minimum are somewhat more intense, and the maximum is slightly shifted to the red, relative to H_2O . The intensity trends suggest that addition of methanol enhances 12-helix population, which mirrors well-established trends for alcohol cosolvents among α -helix-forming conventional peptides;¹⁵ however, strong similarities between NOE data sets for **1** in water and methanol⁶ suggest that the solvent effect on 12-helix population is not large.¹⁶ The similarities among CD data for **1–4** indicate that neither the sulfonamide groups nor the aromatic side chains exert a large effect on the 12-helical CD signature.⁶

We have shown that S-APC residues represent a general strategy for introducing specific side chains at defined positions along the surface of 12-helical β -peptides. The β -peptide 12-helix and the α -helix of conventional peptides have similar geometric features (inner diameter: 3.1 vs 3.2 Å; rise per turn: 5.5 vs 5.4 Å; helix dipole: positive to negative from N-terminus to C-terminus), which suggests that specifically functionalized 12-helices may be able to mimic the structure of α -helical segments of natural proteins. 12-Helical β -peptides may therefore offer a rational approach to development of specific inhibitors of protein–protein interactions that depend on α -helix recognition.¹⁷ β -Peptides containing S-APC and other five-membered-ring-constrained residues are especially interesting from this perspective because of their high conformational stability relative to conventional peptides, which do not form α -helices with fewer than 10–15 residues.^{8,18}

Supporting Information Available: CD data for **1–4** in water and in MeOH; NOE lists for **1** in water and MeOH; summary of Fmoc-S-APC residue synthesis (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA010734R

(11) For leading references, see: Johnson, W. C. *Annu. Rev. Biophys. Biophys. Chem.* **1988**, *17*, 145. Woody, R. W. In *The Peptides*; Academic Press: New York, 1985; Vol. 7, Chapter 2.

(12) Applequist, J.; Bode, K. A.; Appella, D. H.; Christianson, L. A.; Gellman, S. H. *J. Am. Chem. Soc.* **1998**, *120*, 4891.

(13) (a) Abele, S.; Guichard, G.; Seebach, D. *Helv. Chim. Acta* **1998**, *81*, 2141. (b) Gung, B. W.; Zou, D.; Stalcup, A. M.; Cottrell, C. E. *J. Org. Chem.* **1999**, *64*, 2176.

(14) Green, M. M.; Peterson, N. C.; Sato, T.; Teramoto, A.; Cook, R.; Lifson, S. *Science* **1995**, *268*, 1860. Yashima, E.; Maeda, Y.; Okamoto, Y. *J. Am. Chem. Soc.* **1998**, *120*, 8895. Brunsveld, L.; Prince, R. B.; Meijer, E. W.; Moore, J. S. *Org. Lett.* **2000**, *2*, 1525.

(15) (a) Goodman, M.; Verdini, A. S.; Toniolo, C.; Phillips, W. D.; Bovey, F. A. *Proc. Natl. Acad. Sci. U.S.A.* **1969**, *64*, 444. (b) Luo, P.; Baldwin, R. L. *Biochemistry* **1997**, *36*, 8413. (c) Cammers-Goodwin, A.; Allen, T. J.; Oslick, S. L.; McClure, K. F.; Lee, J. H.; Kemp, D. S. *J. Am. Chem. Soc.* **1996**, *118*, 3082. (d) Walgers, R.; Lee, T. C.; Cammers-Goodwin, A. *J. Am. Chem. Soc.* **1998**, *120*, 5073.

(16) Switching from water to methanol exerts a profound effect on β -peptides lacking cyclic backbone constraints, which usually display little or no secondary structure in water: ref 13a and references therein.

(17) For leading references, see: (a) Zutshi, R.; Brickner, M.; Chmielewski, J. *Curr. Opin. Chem. Biol.* **1998**, *2*, 62. (b) Daugherty, D. L.; Gellman, S. H. *J. Am. Chem. Soc.* **1999**, *121*, 4325. (c) Chin, J. W.; Schepartz, A. *J. Am. Chem. Soc.* **2001**, *123*, 2929.

(18) This work was supported by the National Institutes of Health (GM56414) and the Leukemia Society of America. NMR measurements were made in the Department of Chemistry (NIH 1 S10 RR04981) and at the National Magnetic Resonance Facility at Madison (NIH RR02301, RR02781 and RR08438; NSF DMB-8415048 and BIR-9214394; U.S. Department of Agriculture). CD measurements were made in the Biophysics Instrumentation Facility (NSF BIR-9512577).