

ALTERING THE RESIDUES OF RIBONUCLEASE A

R. T. Raines

Department of Biochemistry
University of Wisconsin-Madison
Madison, WI 53706-1569
U. S. A.

ABSTRACT: The cDNA that codes for cow pancreatic ribonuclease A has been cloned and expressed in the yeast *S. cerevisiae*. Site-directed mutagenesis of conserved residues Lys₄₁ and Asp₁₂₁ has been used to probe catalysis by the enzyme. In addition, mutation of Lys₁ to cysteine has allowed for the covalent attachment of a DNA oligonucleotide that confers sequence specificity on ribonuclease A. Finally, a nutritional selection and an activity screen have been developed to distinguish yeast colonies that secrete active ribonuclease A.

The high ribonuclease activity in the pancreas of ruminants has led to the detailed characterization of ribonuclease A (RNase A; EC 3.1.27.5), the predominant ribonuclease in cow pancreas. RNase A has been a widely used subject for protein folding studies; and its structure, its protein chemistry, and the mechanism and energetics of its catalysis are all well-defined (RICHARDS and WYCKOFF, 1971; KARPEISKY and YAKOVLEV, 1981; BLACKBURN and MOORE, 1982; BEINTEMA, 1987). Although RNase A may be better characterized than any other enzyme, it has only recently been subjected to study by molecular biology (STACKHOUSE et al., 1990). Two obstacles have been encountered. First, the





Figure 1. Mechanism of the reaction catalyzed by ribonuclease A. In the transphosphorylation step, His₁₂ is a general base and His₁₁₉ is a general acid. In the hydrolysis step, His₁₁₉ is a general base and His₁₂ is a general acid.

cDNA of RNase A is difficult to clone because the corresponding RNA must be isolated from the pancreas, an organ rich in ribonuclease. Secondly, RNase A is cytotoxic, making the *in vivo* expression of its cDNA problematic.

We have been able to construct a cow pancreas cDNA library from which we have isolated a clone containing the sequence that codes for RNase A. (This sequence is identical to that reported for the RNase A gene by CARSANA et al., 1988). We have been able to express this cDNA in the yeast *S. cerevisiae*, under the control of the repressible acid phosphatase (PHO5) promoter and the glyceraldehyde-3-phosphate



dehydrogenase (GAPDH) transcription terminator (ROSENBERG et al., 1984). In our expression system (Figure 2), RNase A is synthesized as a hybrid with the yeast α -factor leader peptide (BRAKE et al., 1984). The hybrid protein is processed efficiently by endogenous yeast proteases (FULLER et al., 1989), and active RNase A identical to that isolated from cow pancreas is secreted into the media in a quantity (approximately 5 mg/L) adequate for stoichiometric as well as catalytic studies. The RNase A produced is readily purified by cation-exchange chromatography and affinity chromatography on a single-stranded DNA matrix.

Figure 2. Plasmid for the expression of ribonuclease A in yeast.

To understand the relationship between enzyme structure and function and to evaluate the role of particular amino acid side chains in catalysis, it is now common practice to observe the functional consequences of changing active-site residues using recombinant DNA (KNOWLES, 1987). Still, such observations lead to chemical interpretations only when the three-dimensional structure of the enzyme is known. RNase A was first crystallized fifty years ago (KUNITZ, 1939), and structures of both RNase A and the subtilisin-cleaved form, RNase S, and complexes of these with various inhibitors and substrate analogues have been determined by X-ray crystallography and NMR spectroscopy. Equipped with this structural information, we are probing RNase A catalysis using site-directed mutagenesis.

Transition State Stabilization

In 1948 Linus Pauling, in a seminal statement, said: "I think that enzymes are molecules that are complementary in structure to the activated complexes of the

reactions that they catalyze" (PAULING, 1948). Lys₄₁ of RNase A may be an exquisite example of a residue that fulfills Pauling's dictum, since the integral role of Lys₄₁ in RNase A catalysis that is suggested by chemical modification studies (RICHARDS and WYCKOFF, 1971; BLACKBURN and MOORE, 1982) and by its conservation in all 41 sequenced pancreatic ribonucleases (BEINTEMA, 1987) seems to be evident only in the X-ray structure of the complex with uridine vanadate, a transition state analogue (Figure 3)(ALBER et al., 1982). There, the N_ε of Lys₄₁ is hydrogen-bonded to an oxygen atom of the vanadate.




Figure 3. Putative structure of the rate-limiting transition state for the reaction catalyzed by ribonuclease A, and the uridine vanadate analogue of that structure.

To explore the role of Lys₄₁ in RNase A catalysis, we have changed it to arginine, histidine, or methionine and assayed catalysis by these mutant enzymes by following the release of A from UpA (IPATA and FELICIOLI, 1968) and the competitive inhibition of catalysis by uridyl vanadate. Our results indicate that the affinity of a mutant RNase A for the transition state analogue is reflected in its affinity for the rate-limiting transition state (that is, $\text{mutant}(k_{\text{cat}}/K_m)/\text{wild-type}(k_{\text{cat}}/K_m) \approx \text{wild-type}K_i/\text{mutant}K_i$) but not in its affinity for ground states (that is, $\text{wild-type}K_i/\text{mutant}K_i \neq \text{wild-type}K_m/\text{mutant}K_m$). These data are consistent with the side-chain of residue 41 binding exclusively to the rate-limiting transition state. We are proceeding by determining the ability of the mutant enzymes to catalyze each of the half reactions shown in Figure 1.

Catalytic Triad

RNase A may have converged upon a 'catalytic triad' similar to that of the serine proteases. Specifically, Asp₁₂₁ seems to stabilize the tautomer of His₁₁₉ that is appropriate for both the delivery of a proton to the leaving nucleoside and the abstraction of a proton from the water molecule that attacks the cyclic phosphodiester intermediate (shown in Figure 4). [His₁₁₉ and Asp₁₂₁ are conserved in all 41 sequenced pancreatic ribonucleases (BEINTEMA, 1987)]. We are beginning to dissect the putative catalytic triad of RNase A by studying enzymes in which Asp₁₂₁ has been changed to either asparagine (thereby removing the negative charge) or alanine (thereby

removing both the negative charge and the ability to hydrogen bond). Our preliminary studies indicate that both of these mutant RNase A's catalyze the release of A from UpA at the same rate, which is only about 10-fold less than that of the wild-type enzyme!




Figure 4. Catalytic triad of ribonuclease A. The cyclic phosphodiester intermediate is hydrolyzed by Asp₁₂₁, His₁₁₉, and a molecule of water. The side-chain of Asp₁₂₁ is on the surface of the protein.

Hybrid Sequence-Specific Ribonuclease A

We have constructed (as shown in Figure 5) a hybrid RNase A having a covalently-bound DNA oligonucleotide that is complementary to a sequence beginning at the fifth nucleotide downstream from a desired RNA cleavage site. To allow the oligonucleotide to be attached readily and at a spatially appropriate position, we have replaced Lys 1 with cysteine, to yield a fully active ribonuclease with a unique sulfhydryl group. As shown in Figure 6, the sequence-specificity of RNA cleavage of our first-generation hybrid RNase A is improved significantly over that of wild-type RNase A.

Nutritional Selection and Activity Screen for Ribonuclease A

Efficient methods have been developed for introducing definably random base substitutions into specific DNA sequences (HERMES et al., 1989). Such random mutants can be used to discern rapidly those amino acid residues that are critical to a particular screen or selection. We are applying such systematic genetics to RNase A. We have developed a nutritional selection (based on polymeric RNA as the source of uracil for an auxotrophic yeast strain) and an activity screen [based on the acid solubility of monomeric but not polymeric RNA (HOLLOMAN and DEKKER, 1971)] to distinguish yeast clones that secrete active RNase A. We are now searching for mutant enzymes that have a sequence specificity different than that of wild-type RNase A, which cleaves after only after pyrimidine bases.

ACKNOWLEDGEMENTS: We thank R.N. Zuckermann and P.G. Schultz for assaying the specificity of the hybrid site-specific RNase A.




Figure 5. Scheme for the synthesis and reaction of hybrid sequence-specific ribonuclease A.




Figure 6. Autoradiogram of a 20% (w/v) denaturing polyacrylamide gel showing sequence-specific cleavage of a [³²P]-labeled 62-base RNA by our first-generation hybrid RNase A. Lane 1, RNA only; lane 2, ribonuclease U2 digest (A specific); lane 3, hydroxide cleavage ladder; lane 4, RNase A digest; lane 5, hybrid RNase A digest. The arrowhead indicates the cleavage site targeted by the attached DNA oligonucleotide. A14 is residue 14 of the 62-base RNA.

REFERENCES

- ALBER T., GILBERT W.A., PONZI D.R., PETSKO, G.A. (1982) in *Mobility and Function in Proteins and Nucleic Acids* (PORTER R., O'CONNOR M., WHELAN J., eds) Pitman, London p 4-24.
- BEINTEMA J.J. (1987) *Life Chemistry Reports* 4, 333-389.
- BLACKBURN P., MOORE S. (1982) *The Enzymes* 3rd ed, Vol 15, p 317-433.
- BRAKE A.J., MERRYWEATHER J.P., COIT D.G., HEBERLEIN U.A., MASIARZ F.R., MULLENBACH G.T., URDEA M.S., VALENZUELA P., BARR P.J. (1984) *Proc. Natl. Acad. Sci. USA* 81, 4642-4646.
- CARSANA A., CONFALONE E., PALMIERI M., LIBONATI M., FURIA A. (1988) *Nucleic Ac. Res.* 16, 5491-5502.
- FULLER R.S., BRAKE A., THORNER J. (1989) *Proc. Natl. Acad. Sci. USA* 86, 1434-1438.
- HERMES J.D., PAREKH S.M., BLACKLOW S.C., KOSTER H., KNOWLES J.R. (1989) *Gene* 84, 143-151.
- HOLLOMAN W.K., DEKKER C.A. (1971) *Proc. Natl. Acad. Sci. USA* 68, 2241-2245.
- IPATA P.L., FELICOLI R.A. (1968) *FEBS Lett.* 1, 29-31.
- KARPEISKY M.YA., YAKOVLEV, G.I. (1981) *Soviet Scientific Review*, Section D, Vol 2, Harvard Academic Press, New York, p 145-257.
- KNOWLES J.R. (1987) *Science* 236, 1252-1258.
- KUNITZ M. (1939) *Science* 90, 112.
- PAULING L. (1948) *Nature* 161, 707-709.
- RICHARDS F.M, WYCKOFF H.W. (1971) *The Enzymes* 3rd ed, Vol 4, p 647-806.
- ROSENBERG S., BARR P.J., NAJARIAN R.C., HALLEWELL R.A. (1984) *Nature* 312, 77-80.
- STACKHOUSE J., PRESNELL S.R., MCGEEHAN G.M., NAMBIAR K.P., BENNER S.A. (1990) *FEBS Lett.* 262, 104-106.