

# 4-Chloroprolines: Synthesis, Conformational Analysis, and Effect on the Collagen Triple Helix

Matthew D. Shoulders,<sup>1</sup> Ilia A. Guzei,<sup>1</sup> Ronald T. Raines<sup>1,2</sup>

<sup>1</sup> Department of Chemistry, University of Wisconsin–Madison, 1101 University Avenue, Madison, WI 53706-1322

<sup>2</sup> Department of Biochemistry, University of Wisconsin–Madison, 433 Babcock Drive, Madison, WI 53706-1544

Received 15 August 2007; revised 2 October 2007; accepted 5 October 2007

Published online 15 October 2007 in Wiley InterScience (www.interscience.wiley.com). DOI 10.1002/bip.20864

## ABSTRACT:

Collagen is an abundant, triple-helical protein comprising three strands of the repeating sequence: Xaa–Yaa–Gly. (2S)-Proline and (2S,4R)-4-hydroxyproline (Hyp) are common in the primary structure of collagen. Here, we use nonnatural proline derivatives to reveal determinants of collagen stability. Specifically, we report high-yielding syntheses of (2S,4S)-4-chloroproline (clp) and (2S,4R)-4-chloroproline (Clp). We find that the molecular structure of Ac-Clp-OMe in the solid state is virtually identical to that of Ac-Hyp-OMe. In contrast, the conformational properties of Ac-clp-OMe are similar to those of Ac-Pro-OMe. Ac-Clp-OMe has a stronger preference for a trans amide bond than does Ac-Pro-OMe, whereas Ac-clp-OMe has a weaker preference. (Pro–Clp–Gly)<sub>10</sub> forms triple helices that are significantly more stable than those of (Pro–Pro–Gly)<sub>10</sub>. Triple helices of (clp–Pro–Gly)<sub>10</sub> have stability similar to those of (Pro–Pro–Gly)<sub>10</sub>. Unlike (Pro–Clp–Gly)<sub>10</sub> and

(clp–Pro–Gly)<sub>10</sub>, (clp–Clp–Gly)<sub>10</sub> does not form a stable triple helix, presumably due to a deleterious steric interaction between proximal chlorines on different strands. These data, which are consistent with previous work on 4-fluoroprolines and 4-methylprolines, support the importance of stereoelectronic and steric effects in the stability of the collagen triple helix and provide another means to modulate that stability. © 2007 Wiley Periodicals, Inc. *Biopolymers* 89: 443–454, 2008.

**Keywords:** 4-chloroproline; collagen; polyproline II-type helix; proline; prolyl peptide-bond isomerization; triple helix

This article was originally published online as an accepted preprint. The “Published Online” date corresponds to the preprint version. You can request a copy of the preprint by emailing the *Biopolymers* editorial office at [biopolymers@wiley.com](mailto:biopolymers@wiley.com)

## INTRODUCTION

Collagen is the major proteinaceous component of the extracellular matrix (ECM) in vertebrates.<sup>1</sup> The three-dimensional structure of collagen was determined in the 1950s<sup>2–6</sup> to be a right-handed triple helix formed by the parallel coiling of three left-handed polyproline II-type (PPII) strands about a common axis. At least 28 different members of the collagen superfamily of proteins have been discovered to date, as well as a number of other proteins with triple-helical, collagenous domains.<sup>7,8</sup>

In the fibrillar collagens and in most other types of collagen, every third residue is always the smallest of the twenty common amino acids, Gly,<sup>7,9,10</sup> which is required for the tight packing of the triple helix.<sup>11</sup> Often, the amino acid in the Xaa position of the Xaa–Yaa–Gly repeat is Pro and that in

This work is dedicated to the memory of Elkan R. Blout, a pioneer of spectroscopic methods for the structural characterization of proteins and peptides. Professor Blout was among the first to employ collagen-mimetic peptides for the study of native collagen structure, an approach that continues to provide important insights.

Correspondence to: Ronald T. Raines; e-mail: [raines@biochem.wisc.edu](mailto:raines@biochem.wisc.edu)  
Contract grant sponsors: NIH (AR44276, RR13790, P41 RR02301), NSF (BIR-9512577)

© 2007 Wiley Periodicals, Inc.


the Yaa position is (2*S*,4*R*)-4-hydroxyproline (Hyp).<sup>12,13</sup> Hyp is formed by the stereospecific posttranslational hydroxylation of proline residues in the Yaa position by the enzyme prolyl-4-hydroxylase (P4H).<sup>14</sup>

The posttranslational hydroxylation of prolines in the Yaa position is essential for the formation of a stable ECM in animals. Both *Caenorhabditis elegans* and mice lacking P4H experience embryonic morbidity due to weakened collagen superstructures.<sup>15,16</sup> Why is proline hydroxylation essential to the formation of a stable ECM? It is known that collagens containing a high fraction of Hyp in the Yaa position are particularly stable,<sup>17,18</sup> but the physicochemical basis for this stability was unclear.

Determining the basis for the impact of proline hydroxylation on collagen stability is difficult because the high molecular weight and insolubility of native collagen prevent high-resolution structural analysis of the protein in its native form. Elkan Blout and his coworkers were among the first to address this obstacle by employing the reductionist approach of using peptide mimics of collagen, known as collagen-related peptides (CRPs), to reveal fundamental aspects of triple-helix stability.<sup>19–21</sup>

The study of CRPs has led to a series of landmark discoveries regarding the structure and stability of natural collagen. Prockop and coworkers used CRPs to demonstrate that both the stereochemistry and location of Hyp residues are important for its stabilizing effect on collagen triple helices, as illustrated by the  $T_m$  values for triple-helix denaturation listed in Table I.<sup>22–24</sup> CRPs with Hyp in the Yaa position are more stable than those with Pro, but Hyp in the Xaa position prevents triple-helix formation when the Yaa amino acid is Pro.<sup>24,31–35</sup>

In the last decade, we have demonstrated that the stability (or instability) conferred by Hyp derives from the manifesta-



**FIGURE 1** Ring conformations of 4-substituted prolines. The  $C^\gamma$ -endo conformation is favored strongly by stereoelectronic effects when  $R_1 = \text{H}$ ,  $R_2 = \text{F}$  (flp) or  $\text{Cl}$  (clp) and by steric effects when  $R_1 = \text{Me}$  (mep),  $R_2 = \text{H}$ . The  $C^\gamma$ -exo conformation is favored strongly by stereoelectronic effects when  $R_1 = \text{OH}$  (Hyp),  $\text{F}$  (Flp) or  $\text{Cl}$  (Clp),  $R_2 = \text{H}$  and by steric effects when  $R_1 = \text{H}$ ,  $R_2 = \text{Me}$  (Mep).

tion of previously unappreciated stereoelectronic effects. We did so by installing functional groups at  $C^\gamma$  that mediate such effects, as well as reciprocal steric effects (Figure 1).<sup>27,36</sup> For example, replacing Hyp with (2*S*,4*R*)-4-fluoroproline (Flp), which has the native-like stereochemistry, but not (2*S*,4*S*)-4-fluoroproline (flp) results in triple helices with markedly enhanced stability (Table I).<sup>25,29,37</sup> This and other results revealed that a *gauche* effect in Hyp and Flp mandates a  $C^\gamma$ -exo pucker in the pyrrolidine ring,<sup>25,38</sup> which preorganizes the  $\phi$ ,  $\psi$ , and  $\omega$  dihedral angles to those required for triple-helix assembly.<sup>25,39</sup> (2*S*,4*S*)-4-Methylproline (Mep) achieves the same end by manifesting a steric rather than stereoelectronic effect (Figure 1; Table I).<sup>27</sup> (The pyrrolidine ring of proline actually prefers two distinct twist, rather than envelope, conformations.<sup>40</sup> As  $C^\gamma$  experiences a large out-of-plane displacement in these twisted rings, we refer to pyrrolidine ring conformations simply as “ $C^\gamma$ -exo” and “ $C^\gamma$ -endo”.)

The  $C^\gamma$ -endo pucker of proline derivatives in the Xaa position is favorable for triple-helix stability.<sup>26,27,41</sup> This finding was presaged by Zagari and coworkers, who observed that prolines in the Xaa position of high-resolution crystal structures of CRPs nearly always exhibit the  $C^\gamma$ -endo pucker.<sup>42</sup> Consequently, substitution of flp or (2*S*,4*R*)-4-methylproline (mep) for Pro in the Xaa position of CRPs stabilizes triple helices (Table I).<sup>26,27,43,44</sup> Although the  $C^\gamma$ -endo pucker preorganizes the  $\phi$  and  $\psi$  angles to those required for triple-helix assembly, it does not preorganize the  $\omega$  angle because proline derivatives with a  $C^\gamma$ -endo pucker have an enhanced preference for the *cis* peptide bond due to another stereoelectronic effect—an  $n \rightarrow \pi^*$  interaction—described elsewhere,<sup>38,45–48</sup> whereas all peptide bonds in collagen are *trans*. Despite its preference for the  $C^\gamma$ -endo pucker, (2*S*,4*S*)-4-hydroxyproline (hyp) in the Xaa position does not allow for the formation of stable triple helices (Table I),<sup>23</sup> perhaps due to deleterious hydration absent from flp and mep or to idiosyncratic conformational preferences.

**Table I** Effect of 4-Hydroxyproline, 4-Fluoroproline, and 4-Methylproline Diastereomers on the Conformational Stability of Collagen Triple Helices


(Xaa–Yaa–Gly) <sub>n</sub>	$T_m$ (°C)	Ref.
(Pro–Flp–Gly) <sub>7</sub>	45	25
(Pro–Hyp–Gly) <sub>7</sub>	36	25
(flp–Pro–Gly) <sub>7</sub>	33	26
(Pro–Mep–Gly) <sub>7</sub>	29	27
(mep–Pro–Gly) <sub>7</sub>	13	27
(Pro–Pro–Gly) <sub>7</sub>	no helix	28
(Pro–Hyp–Gly) <sub>10</sub>	69	29
(Pro–Pro–Gly) <sub>10</sub>	31–41	29,30
(Hyp–Pro–Gly) <sub>10</sub>	no helix	24
(Pro–hyp–Gly) <sub>10</sub>	no helix	23
(hyp–Pro–Gly) <sub>10</sub>	no helix	23

Thus, steric and stereoelectronic effects can endow great stability on collagen triple helices.<sup>27,36</sup> Beneficial preorganization prescribes that triple helices should be stabilized by proline derivatives in the Xaa position that prefer the C<sup>γ</sup>-endo pucker and by proline derivatives in the Yaa position that prefer the C<sup>γ</sup>-exo pucker. We reasoned that 4-chloro substitution, as in (2*S*,4*S*)-4-chloroprolinone (clp) and (2*S*,4*R*)-4-chloroprolinone (Clp), could control proline ring pucker and modulate collagen stability in much the same manner as does a 4-fluoro substitution (Figure 1).

Chlorine and fluorine have similar physicochemical properties. Organic fluorine has an especially weak propensity to form hydrogen bonds.<sup>49</sup> Hydrogen bonds to organic chlorine (e.g., O—H···Cl—C) can be slightly stronger, but are still relatively weak.<sup>50</sup> Chlorine is the third-most electronegative element after oxygen and fluorine ( $\chi_{\text{O}} = 3.5$ ;  $\chi_{\text{F}} = 4.0$ ;  $\chi_{\text{Cl}} = 3.0$ ).<sup>51</sup> The inductive effect manifested by a chloro group is actually greater than that of a hydroxyl group and similar to that of a fluoro group ( $F_{\text{OH}} = 0.33$ ;  $F_{\text{F}} = 0.45$ ;  $F_{\text{Cl}} = 0.42$ ).<sup>52</sup> The stereoelectronic effect that controls proline ring pucker can be regarded as a hyperconjugative effect,<sup>38</sup> and natural bond orbital analysis suggests that  $\sigma^*_{\text{C-Cl}}$  orbitals have a greater acceptor ability than do the corresponding  $\sigma^*_{\text{C-F}}$  orbitals.<sup>53</sup> Nevertheless, proline ring pucker is controlled by competing stereoelectronic and steric effects,<sup>27,38</sup> and chlorine has a significantly larger covalent radius than does fluorine ( $r_{\text{F}} = 0.64 \text{ \AA}$ ;  $r_{\text{Cl}} = 0.77 \text{ \AA}$ ). Hence, it was not apparent a priori whether the electronegativity or the size of a 4-chloro substitution would dominate its effect on proline ring pucker.

4-Chloroprolinone residues have not been studied in detail in any context. A few chlorinated proline moieties have been found in natural products, all of which are cyclic peptides.<sup>54</sup> Cyclochloritine<sup>55</sup> and astins A–C,<sup>56–59</sup> which are cyclic pentapeptide toxins from *Penicillium islandicum* and *Aster tataricus*, respectively, contain a (2*R*,3*S*,4*R*)-3,4-dichloroprolinone moiety (which is a derivative of L-proline with all three substituents on the same side of the pyrrolidine ring). Likewise, (2*S*,3*S*,4*R*)-3-hydroxy-4-chloroprolinone is a component of astin I.<sup>60</sup> Lee et al. demonstrated that replacing the (2*R*,3*S*,4*R*)-3,4-dichloroprolinone moiety in cyclochloritine with clp decreased its toxicity.<sup>61</sup> Okumura et al. found that clp and especially Clp were incorporated readily into viridogrisein by *Streptomyces griseoviridis* G-89.<sup>62</sup> Mauger and Thomas reported that clp or Clp have different effects on the conformation of actinomycin.<sup>63</sup> Finally, Chiba et al. synthesized very late antigen-4 antagonists containing a clp or Clp residue.<sup>64</sup>

To the best of our knowledge, the effects of 4-chloro substitution on proline ring pucker and peptide-bond isomerization have not been described previously. Likewise, neither



SCHEME 1

clp nor Clp have been incorporated in CRPs or other acyclic peptides. Here, we report on the effect of 4-chloro substitution on the conformation of proline rings, prolyl peptide-bond isomerization, and collagen triple-helix stability.


## RESULTS AND DISCUSSION

### Synthesis of 4-Chloroprolinone Residues

Clp and clp derivatives appropriately protected for peptide synthesis were prepared from the corresponding hydroxyproline derivatives in 97% and 92% overall yield, respectively, via the route shown in Scheme 1. Boc-Hyp-OBn (1) and Boc-hyp-OBn (2) prepared as we described previously<sup>28</sup> were converted to Boc-clp-OBn (3) and Boc-Clp-OBn (4) via an Appel reaction,<sup>65</sup> which had been used previously for the synthesis of 4-chloroprolinone derivatives.<sup>64,66</sup> Subsequent hydrogenolysis of the benzyl group and then exchange of the *N*-Boc protecting group for an *N*-Fmoc protecting group afforded Fmoc-clp-OH (7) and Fmoc-Clp-OH (8).

### Ring Pucker and $K_{\text{trans/cis}}$ of clp and Clp

Compounds of the form Ac-Xaa-OMe are useful model systems for studying proline ring conformation and peptide-bond isomerization, and have been employed both by our



**FIGURE 2** (A) Molecular drawing of crystalline Ac-Clp-OMe (**10**; 50% probability ellipsoids). (B) Conformation of crystalline **10** and Ac-Hyp-OMe<sup>67</sup> depicted with the program PyMOL (Delano Scientific, Palo Alto, CA).

group and by others.<sup>27,38,47,67–69</sup> To determine the effect of 4-chloro substitution on proline ring pucker and peptide-bond isomerization, we prepared Ac-clp-OMe (**9**) and Ac-Clp-OMe (**10**), as shown in Scheme 1. Acidic methanol was used to cleave the *N*-Boc group and introduce the methyl ester.<sup>70</sup> Subsequent treatment with acetyl chloride (<sup>13</sup>C-labeled acetyl chloride was used to aid structural analysis as described in the Materials and Methods section) and *N,N*-dimethylaminopyridine afforded the desired *N*-acetylated target compounds.

Values of  $K_{\text{trans/cis}}$  for molecules of the type Ac-Xaa-OMe correlate closely with proline ring pucker. Proline itself has a slight preference for the *C*<sup>γ</sup>-endo ring pucker, and  $K_{\text{trans/cis}} = 4.6$  in water.<sup>25</sup> Our group has shown that the minimum-energy conformation of 4-substituted derivatives of proline with  $K_{\text{trans/cis}} > 4.6$  is generally a *C*<sup>γ</sup>-exo ring pucker, whereas those with  $K_{\text{trans/cis}} \leq 4.6$  typically prefer a *C*<sup>γ</sup>-endo ring pucker.<sup>38</sup> Using <sup>13</sup>C NMR spectroscopy, we determined that  $K_{\text{trans/cis}} = 5.4$  for Ac-Clp-OMe (**10**) and 2.2 for Ac-clp-OMe (**9**).

We were able to obtain a crystal structure of Ac-Clp-OMe (**10**). Crystalline Ac-Clp-OMe (**10**) displayed the *C*<sup>γ</sup>-exo ring pucker, as expected, and the same overall conformation as we observed previously in the crystal structure of Ac-Hyp-OMe (Figure 2).<sup>67</sup> Notably, the structure of Ac-Clp-OMe (**10**) indicates the presence of a strong  $n \rightarrow \pi^*$  interaction between the amide oxygen and the ester carbonyl, as the  $O'_{i-1} \cdots C'_i$  distance is  $\delta_{\text{BD}} = 2.80$  Å and the  $O'_{i-1} \cdots C'_i=O_i$  angle is  $\tau_{\text{BD}} = 94.1^\circ$ . This  $n \rightarrow \pi^*$  interaction is known to stabilize the *trans* conformation of the amide bond.<sup>38,45–48</sup> Structural parameters for crystalline Ac-Clp-OMe (**10**) and Ac-Hyp-OMe<sup>67</sup> are listed in Table II.

### Synthesis of Chlorinated CRPs

With these results in hand, we suspected that Ac-Clp-OMe (**10**) prefers the *C*<sup>γ</sup>-exo ring pucker and Ac-clp-OMe (**9**) prefers the *C*<sup>γ</sup>-endo ring pucker. Thus, we expected clp to stabilize triple helices in the Xaa position and Clp to stabilize them in the Yaa position.<sup>42</sup> To test this hypothesis, we synthesized (clp-Pro-Gly)<sub>7/10</sub>, (Pro-Clp-Gly)<sub>7/10</sub>, (Clp-Pro-Gly)<sub>10</sub>, and (clp-Clp-Gly)<sub>10</sub>. These CRPs were prepared by segment condensation of the tripeptides Fmoc-clp-Pro-Gly-OH (**18**), Fmoc-Pro-Clp-Gly-OH (**13**), Fmoc-Clp-Pro-Gly-OH (**20**), and Fmoc-clp-Clp-Gly-OH (**15**), respectively, on a solid phase. The Fmoc-protected tripeptides were synthesized as shown in Schemes 2 and 3 using PyBOP<sup>71</sup> and PyBroP<sup>72</sup> to effect the problematic couplings to clp and Clp derivatives.

### Conformational Analysis of clp- and Clp-Containing CRPs


Triple-helical CRPs have a signature circular dichroism (CD) spectrum with a small maximum near 225 nm and a large minimum near 205 nm. This CD spectrum is also characteristic of PPII conformations, but triple helices undergo a cooperative transition upon heating and PPII structures do not. Therefore, we used CD spectroscopy to assess the impact of clp and Clp on triple-helix structure and stability.

The peptides (Pro-Clp-Gly)<sub>7</sub>, (Pro-Clp-Gly)<sub>10</sub>, and (Clp-Pro-Gly)<sub>10</sub> all possess the signature CD spectra of PPII and triple-helical conformations (Figure 3A). Only two of those peptides, (Pro-Clp-Gly)<sub>7</sub> and (Pro-Clp-Gly)<sub>10</sub>, undergo cooperative transitions upon heating with  $T_m$  values of 23 and 52 °C, respectively (Figure 3B; Table III). Thus, our studies indicate that Clp does indeed stabilize triple helices in the Yaa position relative to Pro, although not quite to the same level as Hyp. When Clp is placed in the Xaa position of CRPs, even the long CRP (Clp-Pro-Gly)<sub>10</sub> does not form a triple helix, as indicated by the linear decrease in ellipticity at 225 nm upon heating (Figure 3B and Table III). Thus, both the stereochemistry and position of Clp within CRPs are im-

**Table II** Backbone Dihedral Angles<sup>a</sup> of Crystalline Ac-Clp-OMe (**10**) Derived from X-Ray Diffraction Analysis Compared to Those of Ac-Hyp-OMe<sup>70</sup>

Compound	Ring Pucker	$\phi$	$\psi$	$\omega$	$\delta_{\text{BD}}$ (Å)	$\tau_{\text{BD}}$ (°)
Ac-Clp-OMe ( <b>10</b> )	<i>C</i> <sup>γ</sup> -exo	-56.0	147.5	179.2	2.80	94.1
Ac-Hyp-OMe	<i>C</i> <sup>γ</sup> -exo	-56.9	150.8	178.9	2.82	91.8

<sup>a</sup> For both **10** and Ac-Hyp-OMe, there were two independent molecules in the asymmetric unit of the crystal. Values shown here are the average for these two molecules.




SCHEME 2

portant for its stabilizing effect on triple helices. The self-association of (Pro-Clp-Gly)<sub>10</sub> at 4°C was confirmed by sedimentation equilibrium (see: Supporting Information).


The conformation and stability of clp-containing CRPs was also studied by CD spectroscopy. The effect of clp in the Xaa position on triple-helix stability is similar to that of Pro. Neither (Pro-Pro-Gly)<sub>7</sub> nor (clp-Pro-Gly)<sub>7</sub> forms a stable triple helix. The failure of (clp-Pro-Gly)<sub>7</sub> to form a triple helix is indicated by the linear decrease in ellipticity at 225 nm upon heating (Figure 3D), despite its CD spectrum demonstrating PPII structure at low temperature (Figure 3C). Nonetheless, the longer CRPs (Pro-Pro-Gly)<sub>10</sub> and (clp-Pro-Gly)<sub>10</sub> form triple helices with  $T_m$  values of 31–41<sup>29,30</sup> and 33°C, respectively (Figures 3C and 3D and Table III).

The self-association of (clp-Pro-Gly)<sub>10</sub> at 4°C was confirmed by sedimentation equilibrium (see: Supporting Information).

Two issues must be considered when analyzing the impact of a 4-chloro substitution on proline ring conformation. We have shown that proline ring pucker, and thus  $K_{\text{trans/cis}}$  for the peptide bond, is modulated by reciprocal steric and stereoelectronic effects.<sup>27,38</sup> Chlorine is both more electronegative than methyl and larger than fluorine. Therefore, its impact on proline ring pucker should lie somewhere between the extremes of the small, electron-withdrawing fluorine moiety and the large, electron-donating methyl moiety, both of which strongly enforce proline pucker and thus strongly stabilize (or destabilize) a triple helix. 4-Chloroproline derivatives cannot be expected to endow the same degree of conforma-



SCHEME 3



**FIGURE 3** Conformational analysis of clp- and Clp-containing CRPs. (A, C, and E) CD spectra of peptide solutions (0.2 mM in 50 mM HOAc) at 4°C after incubating at  $\leq 4^\circ\text{C}$  for  $\geq 24$  h. (B, D, and F) Effect of temperature on the molar ellipticity at 225 or 226 nm. Data were recorded at  $3^\circ\text{C}$  intervals after a 5-min equilibration.

tional stability on triple helices as do mep/Mep and flp/Flp or even Hyp, because of the competing effects of chlorine's large size and high electronegativity on proline ring conformation. Thus, the results for triple-helix stability of clp- and Clp-containing CRPs (Table III) are in gratifying agreement with our ideas about collagen strand preorganization.


### Conformational Analysis of (clp-Clp-Gly)<sub>10</sub>

As clp and Clp can be accommodated in the Xaa or Yaa position, respectively, of stable collagen triple helices, it might be expected that the CRP (clp-Clp-Gly)<sub>10</sub> would form a stable triple helix as well. Surprisingly, however, CRPs of the type

(flp-Flp-Gly)<sub>7/10</sub> form significantly less stable triple helices than do the analogous (Pro-Hyp-Gly)<sub>7/10</sub> CRPs.<sup>28,73</sup> Previously, we suggested that this antagonism was due to a deleterious steric interaction between the fluorines of flp and Flp in neighboring strands (Figure 4A).<sup>28</sup> In contrast, Kobayashi and coworkers argued that the instability of triple helices formed from (flp-Flp-Gly) repeats disproved the preorganization theory of triple-helix formation, at least for doubly-substituted CRPs.<sup>73</sup> We have since shown that peptides of the

**Table III** Effect of 4-Chloroproline Diastereomers on the Conformational Stability of Collagen Triple Helices

(Xaa-Yaa-Gly) <sub>n</sub>	T <sub>m</sub> (°C)	Ref.
(Pro-Hyp-Gly) <sub>10</sub>	69	29
(Pro-Clp-Gly) <sub>10</sub>	52	this work
(Pro-Pro-Gly) <sub>10</sub>	31–41	29,30
(clp-Pro-Gly) <sub>10</sub>	33	this work
(flp-Flp-Gly) <sub>10</sub>	30	73
(Pro-Clp-Gly) <sub>7</sub>	23	this work
(clp-Pro-Gly) <sub>7</sub>	<10	this work
(Pro-Pro-Gly) <sub>7</sub>	<10	28
(Clp-Pro-Gly) <sub>10</sub>	<10	this work
(clp-Clp-Gly) <sub>10</sub>	<10	this work



**FIGURE 4** Space-filling models of segments of triple helices constructed from the three-dimensional structure of a (Pro-Hyp-Gly)<sub>n</sub> triple helix (PDB entry 1CAG<sup>11</sup>) by replacing the H or OH on Pro and Hyp with F or Cl, respectively, using the program SYBYL (Tripos, St. Louis, MO) and depicting the images with the program PyMOL. (A) Segment of a (flp-Flp-Gly)<sub>n</sub> triple helix ( $r_{\text{F}\dots\text{F}} = 2.4 \text{ \AA}$ ).<sup>28</sup> (B) Segment of a (clp-Clp-Gly)<sub>n</sub> triple helix ( $r_{\text{Cl}\dots\text{Cl}} = 1.9 \text{ \AA}$ ).

form (mep-Mep-Gly)<sub>7</sub> are more stable than both of the mono-substituted variants (mep-Pro-Gly)<sub>7</sub> and (Pro-Mep-Gly)<sub>7</sub>,<sup>27</sup> substantiating the importance of preorganization in doubly-substituted CRPs. We sought to probe deleterious steric interactions in doubly-substituted CRPs again, now with chlorinated CRPs.

We found that (clp-Clp-Gly)<sub>10</sub> triple helices do not form, even at low temperatures. The (clp-Clp-Gly)<sub>10</sub> CRP does possess a PPII-type structure at low temperature (Figure 3E), but does not fold into a stable triple helix. This instability is demonstrated by the absence of a cooperative transition at 225 nm upon heating (Figure 3F) and by sedimentation equilibrium, which showed no self-assembly at low temperature (see: Supporting Information). In contrast, (flp-Flp-Gly)<sub>10</sub> triple helices have a  $T_m$  of 30°C.<sup>73</sup> These findings suggest that the large size of chlorine relative to fluorine exacerbates the deleterious steric interaction between neighboring strands, as depicted in Figure 4B. 4-Chloroproline derivatives are likely to have greater conformational flexibility than 4-fluoroproline derivatives, so if “locking” the proline rings into a particular deleterious conformation were responsible for the instability of (flp-Flp-Gly)<sub>10</sub>,<sup>73</sup> then (clp-Clp-Gly)<sub>10</sub> triple helices should be stable. They are not, supporting our hypothesis<sup>28</sup> rather than that of Kobayashi and coworkers.<sup>73</sup>

## CONCLUSIONS

Clp has a strong preference for the C<sup>γ</sup>-exo ring pucker and the *trans* amide bond, whereas clp prefers the C<sup>γ</sup>-endo ring pucker and has a lower proportion of the *trans* amide bond. These results could explain the contrasting effects of clp and Clp on the biosynthesis and activity of natural products.<sup>61–63</sup> They further suggest that clp has an appropriate conformation to stabilize polyproline I-type strands whereas Clp should stabilize PPII strands, as was found previously for flp and Flp.<sup>74,75</sup> Additionally, we have demonstrated that Clp in the Yaa position of CRPs confers nearly the same degree of stability to triple helices as does Hyp, but it prevents triple-helix formation when placed in the Xaa position. In contrast, clp behaves much like Pro in a triple-helical context, as (clp-Pro-Gly)<sub>10</sub> and (Pro-Pro-Gly)<sub>10</sub> triple helices have similar thermal stabilities. Triple helices do not form from the doubly-substituted CRP (clp-Clp-Gly)<sub>10</sub>, in contrast to (Pro-Pro-Gly)<sub>10</sub> and (flp-Flp-Gly)<sub>10</sub> CRPs. This instability is likely due to a strongly deleterious steric interaction between the chlorines in neighboring strands. An interesting attribute of clp- and Clp-containing triple helices is that they could be modified covalently by the S<sub>N</sub>2 attack of nucleophiles at the halogenated carbon. Finally, we note the similarity of the active site and reactivity of mammalian P4H, the enzyme re-

sponsible for hydroxylation of Pro residues in natural collagen,<sup>76</sup> and the nonhaem iron halogenase SyrB2,<sup>77</sup> an enzyme responsible for the chlorination of unactivated carbon atoms.<sup>78</sup> As Clp confers thermal stability on collagen triple helices at a level similar to that of Hyp, it is possible that organisms evolving in an environment rich in chloride ions could evolve stable chlorinated collagens analogous to the hydroxylated collagens found in modern animals. The impact of chlorination rather than hydroxylation on the supramolecular structure of native collagen is a subject for future investigation.

## MATERIALS AND METHODS

### General

Commercial chemicals were of reagent grade or better, and were used without further purification. Anhydrous DMF and CH<sub>2</sub>Cl<sub>2</sub> were obtained from CYCLE-TAINER<sup>®</sup> solvent delivery systems (J. T. Baker, Phillipsburg, NJ). Other anhydrous solvents were obtained in septum-sealed bottles. In all reactions using anhydrous solvents, glassware was either oven- or flame-dried. “NaHCO<sub>3</sub>(aq)” and “brine” (i.e., NaCl) refer to saturated aqueous solutions unless specified otherwise. Flash chromatography was performed with columns of silica gel 60, 230–400 mesh (Silicycle, Québec City, Canada). HPLC was performed with gradients of solvent A (0.1% v/v TFA in water) and solvent B (0.1% v/v TFA in acetonitrile), as indicated.

The term “concentrated under reduced pressure” refers to the removal of solvents and other volatile materials using a rotary evaporator at water aspirator pressure (<20 torr) while maintaining the water-bath temperature below 50°C. Residual solvent was removed from samples at high vacuum (<0.1 torr). The term “high vacuum” refers to vacuum achieved by a mechanical belt-drive oil pump.

NMR spectra were acquired with a Bruker DMX-400 Avance spectrometer unless specified otherwise (<sup>1</sup>H, 400 MHz; <sup>13</sup>C, 100.6 MHz) at the National Magnetic Resonance Facility at Madison. NMR spectra were obtained at ambient temperatures on samples dissolved in CDCl<sub>3</sub> or MeOH-*d*<sub>4</sub>. Coupling constants *J* are provided in Hertz. Compounds with a carbamate protecting group (e.g., Boc or Fmoc) exist as mixtures of *Z* and *E* isomers that do not interconvert on the NMR time scale at ambient temperatures. Accordingly, these compounds exhibit two sets of NMR signals.

Mass spectrometry was performed with either a Micromass LCT (electrospray ionization, ESI) in the Mass Spectrometry Facility in the Department of Chemistry or an Applied Biosystems Voyager DE-Pro (matrix-assisted laser desorption/ionization, MALDI) mass spectrometer in the University of Wisconsin Biophysics Instrumentation Facility.

### Synthesis of Boc-clp-OBn (3) and Boc-Clp-OBn (4)

**General Protocol.** The appropriate protected 4-hydroxyproline derivative **1** or **2** (33.8 g, 105 mmol) was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (115 mL) under Ar(g) and cooled to 0°C. Triphenylphosphine (49.6 g, 189 mmol) and then CCl<sub>4</sub> (153.8 g, 96 mL) were

added. The reaction mixture was stirred at 0°C for 3 h, then heated to 35°C for 2 h, and then cooled to rt and stirred for 45 min. The reaction mixture was concentrated under reduced pressure, and the residue was purified by flash chromatography over silica gel (20% v/v EtOAc in hexane).

***N*-tert-Butyloxycarbonyl-(2*S*,4*S*)-4-chloroproline Benzyl Ester (3).** Boc-clp-OBn (**3**) was obtained in 92% yield as a colorless liquid. <sup>1</sup>H NMR δ: 1.34 and 1.46 (s, 9H), 2.38 (dt, *J* = 5.2, 13.9, 1H), 2.64–2.78 (m, 1H), 3.65 (td, *J* = 4.7, 12.7, 1H), 3.88–4.01 (m, 1H), 4.37 and 4.51 (m, 2H), 5.06–5.32 (m, 2H), 7.29–7.40 (m, 5H); <sup>13</sup>C NMR δ: 28.2, 28.4, 39.5, 40.4, 53.7, 54.7, 55.0, 55.4, 57.7, 58.0, 67.1, 80.6, 128.2, 128.5, 128.6, 128.6, 135.4, 135.6, 153.3, 153.8, 171.3, 171.6. ESI-MS (*m/z*): [2M + Na]<sup>+</sup> Calcd for C<sub>34</sub>H<sub>44</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>8</sub>Na 701.2; found 701.9 (2 Cl).

***N*-tert-Butyloxycarbonyl-(2*S*,4*R*)-4-chloroproline Benzyl Ester (4).** Boc-Clp-OBn (**4**) was obtained in 97% yield as a colorless liquid. <sup>1</sup>H NMR δ: 1.36 and 1.47 (s, 9H), 2.28–2.57 (m, 2H), 3.64–3.90 (m, 2H), 4.47 (m, 1H), 4.50 and 4.59 (t, *J* = 7.6, 1H), 5.08–5.31 (m, 2H), 7.35 (m, 5H); <sup>13</sup>C NMR (125 MHz) δ: 28.3, 28.5, 39.8, 40.8, 55.3, 55.5, 55.6, 55.7, 57.8, 58.0, 67.2, 80.8, 80.9, 128.3, 128.5, 128.6, 128.7, 128.8, 135.4, 135.7, 153.7, 154.2, 172.1, 172.4; HRMS-ESI (*m/z*): [2M + Na]<sup>+</sup> Calcd for C<sub>34</sub>H<sub>44</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>8</sub>Na 701.2372; found 701.2343 (2 Cl).

### Synthesis of Boc-clp-OH (5) and Boc-Clp-OH (6)

**General Protocol.** MeOH (400 mL) was added carefully to a mixture of the appropriate 4-chloroproline derivative **3** or **4** (31.8 g, 93.6 mmol) and Pd/C (10% w/w, 10.4 g) under Ar(g), and the resulting black suspension was stirred under H<sub>2</sub>(g) for 23 h. The suspension was filtered through a pad of Celite and concentrated under reduced pressure.

***N*-tert-Butyloxycarbonyl-(2*S*,4*S*)-4-chloroproline (5).** Boc-clp-OH (**5**) was obtained in quantitative yield as a white solid. <sup>1</sup>H NMR δ: 1.44 and 1.49 (s, 9H), 2.37–2.85 (m, 2H), 3.56–3.72 (m, 1H), 3.84–4.04 (m, 1H), 4.32–4.51 (m, 2H), 8.22 (bs, 1H); <sup>13</sup>C NMR (125 MHz) δ: 28.3, 28.4, 38.8, 40.3, 53.6, 54.5, 54.9, 55.7, 57.9, 81.3, 81.8, 153.6, 155.3, 175.2, 177.2; HRMS-ESI (*m/z*): [M – H]<sup>–</sup> Calcd for C<sub>10</sub>H<sub>15</sub>ClNO<sub>2</sub> 248.0690; found 248.0694 (1 Cl).

***N*-tert-Butyloxycarbonyl-(2*S*,4*R*)-4-chloroproline (6).** Boc-Clp-OH (**6**) was obtained in quantitative yield as a white solid. <sup>1</sup>H NMR δ: 1.44 and 1.50 (s, 9H), 2.38–2.72 (m, 2H), 3.70–3.83 (m, 2H), 4.49 (m, 1H), 4.57 (t, *J* = 7.2, 1H); <sup>13</sup>C NMR (125 MHz) δ: 28.3, 28.4, 38.9, 40.7, 55.3, 55.4, 55.9, 57.8, 58.0, 81.3, 82.5, 153.6, 156.4, 174.4, 177.9; HRMS-ESI (*m/z*): [M – H]<sup>–</sup> Calcd for C<sub>10</sub>H<sub>15</sub>ClNO<sub>2</sub> 248.0690; found 248.0680 (1 Cl).

### Synthesis of Fmoc-clp-OH (7) and Fmoc-Clp-OH (8)

**General Protocol.** The appropriate 4-chloroproline derivative **5** or **6** (22.6 g, 90.3 mmol) was dissolved in 4 N HCl in dioxane (550 mL) under Ar(g) and stirred for 3 h. The resulting solution was concentrated under reduced pressure. The free amine was dissolved in 10% w/v NaHCO<sub>3</sub>(aq) (375 mL). A solution of Fmoc-OSu (33.5 g, 99.3 mmol) in dioxane (600 mL) was added, and the resulting white suspension was stirred for 19 h. The reaction mixture was concentrated under reduced pressure, and the aqueous solution was diluted with water (300 mL) and washed with ether (3 × 500 mL). The

aqueous layer was acidified to pH 1.5 with 12N HCl, extracted with ether (3 × 500 mL), dried over anhydrous MgSO<sub>4</sub>(s), and concentrated under reduced pressure.

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*S*)-4-chloroproline (7).** Fmoc-clp-OH (**7**) was obtained in quantitative yield as a white solid. <sup>1</sup>H NMR (500 MHz) δ: 2.42–2.77 (m, 2H), 3.63–3.76 (m, 1H), 3.85–4.04 (m, 1H), 4.13–4.62 (m, 5H), 7.27–7.45 (m, 4H), 7.50–7.62 (m, 2H), 7.69–7.80 (m, 2H); <sup>13</sup>C NMR (125 MHz) δ: 38.7, 40.4, 47.2, 53.8, 54.6, 55.7, 57.4, 58.2, 67.8, 68.3, 120.2, 125.0, 125.1, 127.3, 127.9, 128.0, 141.5, 143.6, 143.8, 154.3, 155.5, 174.6, 176.0; HRMS-ESI (*m/z*): [M + Na]<sup>+</sup> Calcd for C<sub>20</sub>H<sub>18</sub>ClNO<sub>4</sub>Na 394.0822; found 394.0828 (1 Cl).

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*R*)-4-chloroproline (8).** Fmoc-Clp-OH (**8**) was obtained in quantitative yield as a white solid. <sup>1</sup>H NMR (500 MHz) δ: 2.35–2.68 (m, 2H), 3.83 (m, 2H), 4.11–4.67 (m, 5H), 7.13 (bs, 1H), 7.23–7.45 (m, 4H), 7.49–7.62 (m, 2H), 7.74 (m, 2H); <sup>13</sup>C NMR (125 MHz) δ: 39.3, 40.8, 47.2, 47.3, 55.0, 55.5, 55.7, 56.0, 57.3, 58.0, 68.0, 68.3, 120.1, 120.2, 124.9, 125.1, 127.2, 127.3, 127.8, 128.0, 141.4, 143.6, 146.7, 144.0, 154.5, 155.8, 175.2, 176.8; HRMS-ESI (*m/z*): [M + Na]<sup>+</sup> Calcd for C<sub>20</sub>H<sub>18</sub>ClNO<sub>4</sub>Na 394.0822; found 394.0812 (1 Cl).

### Synthesis of Ac-clp-OMe (9) and Ac-Clp-OMe (10)

**General Protocol.** The appropriate 4-chloroproline derivative **5** or **6** (107 mg, 0.4 mmol) was dissolved in anhydrous MeOH (8 mL) and cooled to 0°C. Acetyl chloride (8 mL) was added dropwise and the resulting solution was stirred for 7 h at rt. The solution was concentrated under reduced pressure and dried overnight under high vacuum. The residue was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under Ar(g). *N,N*-dimethylaminopyridine (397 mg, 3.3 mmol) and acetyl chloride (250 mg, 3.1 mmol) were added, and the resulting solution was stirred for 22 h. The reaction mixture was concentrated under reduced pressure, and the residue was dissolved in 10% w/v aqueous citric acid (40 mL). The aqueous solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 × 75 mL), which was then dried over anhydrous MgSO<sub>4</sub>(s), and concentrated under reduced pressure. The residue was purified by flash chromatography over silica gel (10% v/v hexane in EtOAc to elute byproducts and then 15% v/v MeOH in EtOAc to elute product).

***N*-(Acetyl)-(2*S*,4*S*)-4-chloroproline Methyl Ester (9).** Ac-clp-OMe (**9**) was obtained in 82% yield as a fragrant, colorless oil. <sup>1</sup>H NMR δ: 2.02 and 2.08 (2 s, 3H), 2.30–2.38 (m, 0.7H), 2.62–2.79 (m, 1.3H), 3.73 and 3.78 (2 s, 3H), 3.72–3.82 (m, 1H), 3.97–4.06 (m, 1H), 4.38–4.49 (m, 1.3H), 4.58 (dd, *J* = 5.2, 8.8, 0.7H); <sup>13</sup>C NMR δ: 22.1, 22.4, 39.0, 41.1, 52.6, 52.9, 54.2, 54.5, 55.8, 56.2, 57.4, 58.8, 169.2, 170.0, 171.4, 171.4; HRMS-ESI (*m/z*): [M + H]<sup>+</sup> Calcd for C<sub>8</sub>H<sub>13</sub>NO<sub>3</sub> 205.0584; found 205.0582 (1 Cl).

***N*-(Acetyl)-(2*S*,4*R*)-4-chloroproline Methyl Ester (10).** Ac-Clp-OMe (**10**) was obtained in 76% yield as a fragrant, colorless oil. <sup>1</sup>H NMR δ: 1.99 and 2.08 (2 s, 3H), 2.31–2.69 (m, 2H), 3.74 and 3.78 (2 s, 3H), 3.69–3.85 (m, 1H), 4.03 (dd, *J* = 5.0, 11.4, 1H), 4.45–4.67 (m, 2H); <sup>13</sup>C NMR δ: 21.7, 22.3, 39.3, 41.4, 52.6, 53.0, 54.4, 55.2, 55.7, 56.7, 57.4, 58.6, 169.4, 170.0, 172.1, 172.4; HRMS-ESI (*m/z*): [M + Na]<sup>+</sup> Calcd for C<sub>8</sub>H<sub>12</sub>NO<sub>3</sub>Na 205.0403; found 205.0403 (1 Cl).



### ***N*-tert-Butyloxycarbonyl-(2*S*,4*R*)-4-chloropropyl-glycine Benzyl Ester (11)**

Boc-Clp-OH (**6**) (23.3 g, 93.2 mmol), glycine benzyl ester tosylate (40.9 g, 121.2 mmol), and PyBOP (48.5 g, 93.2 mmol) were dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (400 mL) under Ar(g). DIEA (30.1 g, 233 mmol) was added slowly, and the resulting solution was stirred for 16 h. The reaction mixture was washed with 10% w/v aqueous citric acid (2 × 1.0 L), dried over anhydrous MgSO<sub>4</sub>(s), and concentrated under reduced pressure. The crude oil was purified by flash chromatography over silica gel (gradient: 33% v/v EtOAc in hexane to 50% v/v EtOAc in hexane) to afford Boc-Clp-Gly-OBn (**11**) (29.8 g, 75.1 mmol, 81%) as a colorless, sticky paste. <sup>1</sup>H NMR δ: 1.47 (s, 9H), 2.25–2.80 (m, 2H), 3.70 and 3.92 (m, 2H), 4.01–4.16 (m, 2H), 4.41–4.59 (m, 2H), 5.18 (s, 2H), 6.61 (m, 0.3H), 7.36 (m, 5H); <sup>13</sup>C NMR δ: 28.4, 38.2, 41.0, 41.3, 41.6, 55.7, 55.9, 58.6, 59.8, 67.3, 67.5, 81.6, 128.5, 128.7, 128.8, 135.3, 157.1, 166.8, 169.5, 171.3; HRMS–ESI (*m/z*): [M + Na]<sup>+</sup> Calcd for C<sub>19</sub>H<sub>25</sub>ClN<sub>2</sub>O<sub>5</sub>Na, 419.1350; found, 419.1335 (1 Cl).

### ***N*-9-Fluorenylmethoxycarbonyl-(2*S*)-prolyl-(2*S*,4*R*)-4-chloropropyl-glycine Benzyl Ester (12)**

Boc-Clp-Gly-OBn (**11**) (1.00 g, 2.5 mmol) was dissolved in 4N HCl in dioxane (30 mL) under Ar(g) and stirred for 2 h. The resulting solution was concentrated under reduced pressure and the residue dissolved in anhydrous DMF (50 mL) under Ar(g). DIEA (1.20 g, 9.3 mmol) was added, followed by Fmoc-Pro-OPfp (2.67 g, 5.3 mmol). The solution was stirred for 18 h and then concentrated by rotary evaporation under high vacuum. Flash chromatography over silica gel (gradient: 20% v/v EtOAc in hexane to 30% v/v EtOAc in hexane) afforded **12** (1.10 g, 1.8 mmol, ~71%) as a white solid containing an impurity that was removed after the subsequent step.

### **PyBroP Couplings**

**General Protocol.** The appropriate Boc-Xaa-Gly-OBn derivative (**11**) or (**16**)<sup>79</sup> (52.5 mmol) was dissolved in 4 N HCl in dioxane (700 mL) under Ar(g) and stirred for 2 h. The resulting solution was concentrated under reduced pressure, dried under high vacuum, and the residue dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (800 mL) under Ar(g). The appropriate 4-chloroprolines derivative (**7**) or (**8**) (52.5 mmol) was added, and the resulting solution was cooled to 0°C. PyBroP (24.5 g, 52.5 mmol) and DIEA (23.8 g, 184 mmol) were added. The resulting solution was allowed to warm slowly to room temperature and then stirred for 18 h. The reaction mixture was washed with 10% w/v aqueous citric acid (1.0 L), NaHCO<sub>3</sub>(aq) (1.0 L), and brine (1.0 L). The organic layer was dried over anhydrous MgSO<sub>4</sub>(s) and concentrated under reduced pressure. The residue was purified by flash chromatography over silica gel (gradient: 30% v/v EtOAc in hexane to 90% v/v EtOAc in hexane).

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*S*)-4-chloropropyl-(2*S*,4*R*)-4-chloro prolyl-glycine Benzyl Ester (14).** Fmoc-clp-Clp-Gly-OBn (**14**) was obtained in 67% yield as a white solid. HRMS–ESI (*m/z*): [M + Na]<sup>+</sup> Calcd for C<sub>34</sub>H<sub>33</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>6</sub>Na, 672.1644; found, 672.1646 (2 Cl).

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*S*)-4-chloropropyl-(2*S*)-prolyl-glycine Benzyl Ester (17).** Fmoc-clp-Pro-Gly-OBn (**17**) was

obtained in 82% yield as a white solid. HRMS–ESI (*m/z*): [M + Na]<sup>+</sup> Calcd for C<sub>34</sub>H<sub>34</sub>ClN<sub>3</sub>O<sub>6</sub>Na, 638.2034; found, 638.2039 (1 Cl).

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*R*)-4-chloropropyl-(2*S*)-prolyl-glycine Benzyl Ester (19).** Fmoc-Clp-Pro-Gly-OBn (**19**) was obtained in 72% yield as a white solid. HRMS–ESI (*m/z*): [M + Na]<sup>+</sup> calcd for C<sub>34</sub>H<sub>34</sub>ClN<sub>3</sub>O<sub>6</sub>Na, 638.2034; found, 638.2009 (1 Cl).

### **Hydrogenolysis of Benzyl Groups on Protected Tripeptides**

**General Protocol.** MeOH (400 mL) was added carefully to a mixture of the appropriate Fmoc-protected tripeptide benzyl ester (**12**), (**14**), (**17**), or (**19**) (34.6 mmol) and Pd/C (10% w/w, 9.6 g) under Ar(g). The resulting black suspension was stirred under H<sub>2</sub>(g) for ~4 h. Careful monitoring by TLC was necessary to prevent hydrogenolysis of the Fmoc group. The suspension was filtered through a pad of Celite and concentrated under reduced pressure. The crude product was purified by flash chromatography over silica gel (EtOAc to elute byproducts, then 12% v/v MeOH in CH<sub>2</sub>Cl<sub>2</sub> containing 0.1% v/v formic acid). The fractions containing the reaction product were concentrated under reduced pressure and the formic acid was removed by dissolving the residue in 10% v/v MeOH in toluene and concentrating under reduced pressure.

***N*-9-Fluorenylmethoxycarbonyl-(2*S*)-prolyl-(2*S*,4*R*)-4-chloropropyl-glycine (13).** Fmoc-Pro-Clp-Gly-OH (**13**) was obtained in 67% yield as a white solid. <sup>13</sup>C NMR (125 MHz, MeOH-*d*<sub>4</sub>) δ: 24.1, 25.2, 30.1, 31.0, 40.1, 42.1, 47.9, 56.7, 57.0, 57.3, 57.4, 59.3, 59.7, 59.9, 60.2, 60.3, 68.4, 68.7, 120.9, 126.1, 126.2, 126.3, 128.2, 128.4, 128.8, 142.5, 142.6, 145.0, 145.1, 145.4, 145.5, 156.2, 156.6, 173.3, 173.3, 173.5, 173.7; HRMS–ESI (*m/z*): [M – H]<sup>–</sup> Calcd for C<sub>27</sub>H<sub>27</sub>ClN<sub>3</sub>O<sub>6</sub>, 524.1589; found, 524.1586 (1 Cl).

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*S*)-4-chloropropyl-(2*S*,4*R*)-4-chloropropyl-glycine (15).** Fmoc-clp-Clp-Gly-OH (**15**) was obtained in 66% yield as a white solid. <sup>13</sup>C NMR (125 MHz, MeOH-*d*<sub>4</sub>) δ: 39.6, 40.0, 40.1, 40.5, 41.9, 53.2, 53.6, 55.6, 56.1, 56.8, 57.1, 57.4, 58.5, 58.8, 60.3, 60.4, 68.8, 68.9, 120.9, 126.1, 126.3, 128.2, 128.4, 128.7, 128.7, 142.5, 142.5, 142.6, 142.7, 144.9, 145.0, 145.3, 145.4, 156.7, 156.0, 172.1, 172.2, 172.6, 173.4, 173.6; HRMS–ESI (*m/z*): [M + Na]<sup>+</sup> Calcd for C<sub>27</sub>H<sub>27</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>6</sub>Na, 582.1175; found, 582.1157 (2 Cl).

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*S*)-4-chloropropyl-(2*S*)-prolyl-glycine (18).** Fmoc-clp-Pro-Gly-OH (**18**) was obtained in 57% yield as a white solid. <sup>13</sup>C NMR (125 MHz, MeOH-*d*<sub>4</sub>) δ: 24.1, 25.7, 25.9, 28.6, 30.1, 30.3, 37.8, 40.3, 41.8, 46.5, 53.0, 53.5, 55.2, 55.5, 56.0, 56.3, 58.3, 58.7, 59.8, 61.3, 61.6, 67.7, 68.9, 120.9, 125.7, 125.9, 126.1, 128.2, 128.9, 142.5, 142.6, 144.9, 145.0, 145.3, 145.5, 155.6, 155.9, 167.7, 169.2, 171.9, 172.1, 172.6, 174.4, 174.6; HRMS–ESI (*m/z*): [M + Na]<sup>+</sup> Calcd C<sub>27</sub>H<sub>28</sub>ClN<sub>3</sub>O<sub>6</sub>Na, 548.1564; found, 548.1561 (1 Cl).

***N*-9-Fluorenylmethoxycarbonyl-(2*S*,4*R*)-4-chloropropyl-(2*S*)-prolyl-glycine (20).** Fmoc-Clp-Pro-Gly-OH (**20**) was obtained in 58% yield as a white solid. <sup>13</sup>C NMR (125 MHz, MeOH-*d*<sub>4</sub>) δ: 24.2, 25.7, 25.9, 28.6, 30.1, 30.4, 39.1, 40.4, 41.0, 42.1, 46.3, 47.9, 56.1, 56.8, 57.1, 57.3, 57.4, 57.7, 58.0, 58.2, 59.9, 61.4, 61.7, 67.7, 68.9, 121.0, 125.7, 125.9, 126.1, 128.2, 128.3, 128.9, 142.6, 142.7, 144.8, 145.0, 145.2, 145.5, 156.1, 156.4, 168.0, 168.6, 172.2, 172.4, 174.2, 174.5; HRMS–ESI (*m/z*): [M + Na]<sup>+</sup> Calcd C<sub>27</sub>H<sub>28</sub>ClN<sub>3</sub>O<sub>6</sub>Na, 548.1564; found, 548.1578 (1 Cl).

### Measurement of $K_{\text{trans/cis}}$ Values of Ac-clp-OMe (9) and Ac-Clp-OMe (10)

$^{13}\text{C}$ -labeled versions of Ac-clp-OMe (9) and Ac-Clp-OMe (10) (5–10 mg) were dissolved in  $\text{D}_2\text{O}$  with sufficient  $\text{MeOH-}d_4$  added to solubilize the compound (<20% of total volume). The  $^{13}\text{C}$  NMR spectra were recorded using an inverse-gated decoupling pulse program with a relaxation delay of 100 s and a pulse width of 10  $\mu\text{s}$ . The spectral baselines were corrected and peaks corresponding to the labeled carbon were integrated. Values of  $K_{\text{trans/cis}}$  were determined by the relative areas of the trans and cis peaks for the labeled carbons.

### General Protocol for Attachment of Fmoc-Pro-Clp-Gly-OH (13), Fmoc-clp-Clp-Gly-OH (15), Fmoc-clp-Pro-Gly-OH (18), and Fmoc-Clp-Pro-Gly-OH (20) to 2-Chlorotrityl Resin

Under  $\text{Ar(g)}$ , 23 mg of 2-chlorotrityl resin (loading: 1.6 mmol/g) was swelled in anhydrous  $\text{CH}_2\text{Cl}_2$  (0.7 mL) for 5 min. A solution of the appropriate Fmoc-protected tripeptide (13), (15), (18), or (20) (0.034 mmol) and DIEA (17 mg, 0.13 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (0.7 mL) was added by syringe. Additional anhydrous  $\text{CH}_2\text{Cl}_2$  (0.5 mL) was used to ensure complete transfer. After 2 h, anhydrous MeOH (0.2 mL) was added to cap any remaining active sites on the resin. The resin-bound peptide was isolated by gravity filtration, washed with several portions of anhydrous  $\text{CH}_2\text{Cl}_2$  (~25 mL), and dried under high vacuum. Loadings were measured by an Fmoc-deprotection ultraviolet spectroscopy assay to be 0.44 mmol/g for (13), 0.55 mmol/g for (15), 0.35 mmol/g for (18), and 0.48 mmol/g for (20).

### Synthesis of (Pro-Clp-Gly) $_7$ , (clp-Pro-Gly) $_7$ , (Pro-Clp-Gly) $_{10}$ , (clp-Pro-Gly) $_{10}$ , (clp-Clp-Gly) $_{10}$ , and (Clp-Pro-Gly) $_{10}$

The two 21-mer peptides and the four 30-mer peptides were synthesized by segment condensation of their corresponding Fmoc-tripeptides (13, 15, 18, and 20) on a solid phase using an Applied Biosystems Synergy 432A Peptide Synthesizer at the University of Wisconsin–Madison Biotechnology Center. The first trimer was loaded onto the resin as described above. Fmoc-deprotection was achieved by treatment with 20% (v/v) piperidine in DMF. The trimers (3 equiv) were converted to active esters by treatment with HBTU, DIEA, and HOBt. Extended couplings (30–60 min) were employed at room temperature.

Peptides were cleaved from the resin in 95:3:2 TFA:triisopropylsilane: $\text{H}_2\text{O}$  (1.5 mL), precipitated from *t*-butylmethylether at 0°C, isolated by centrifugation, and lyophilized. Semipreparative HPLC was used to purify the peptides (Pro-Clp-Gly) $_7$  (Dynamax C-18 column, gradient: 10% B to 65% B over 50 min), (clp-Pro-Gly) $_7$  (Dynamax C-18 column, gradient: 10% B to 65% B over 50 min), (Pro-Clp-Gly) $_{10}$  (Zorbax C-8 column, gradient: 10% B to 65% B over 60 min), (clp-Pro-Gly) $_{10}$  (Zorbax C-8 column, gradient: 10% B to 90% B over 60 min), (clp-Clp-Gly) $_{10}$  (Zorbax C-8 column, gradient: 10% B to 90% B over 60 min), and (Clp-Pro-Gly) $_{10}$  (Zorbax C-8 column, gradient 10% B to 65% B over 60 min). All six peptides were >90% pure by analytical HPLC and MALDI–TOF

mass spectrometry ( $m/z$ )  $[\text{M} + \text{H}]^+$  Calcd for  $\text{C}_{84}\text{H}_{122}\text{Cl}_7\text{N}_{21}\text{O}_{22}$  2020; found 2019 for (Pro-Clp-Gly) $_7$ ;  $[\text{M} + \text{Na}]^+$  Calcd for  $\text{C}_{84}\text{H}_{121}\text{Cl}_7\text{N}_{21}\text{NaO}_{22}$  2044; found 2041 for (clp-Pro-Gly) $_7$ ;  $[\text{M} + \text{Na}]^+$  Calcd for  $\text{C}_{120}\text{H}_{162}\text{Cl}_{10}\text{N}_{30}\text{O}_{31}\text{Na}$  2900; found 2900 for (Pro-Clp-Gly) $_{10}$ , 2897 for (clp-Pro-Gly) $_{10}$ , and 2899 (Clp-Pro-Gly) $_{10}$ ;  $[\text{M} + \text{Na}]^+$  Calcd for  $\text{C}_{120}\text{H}_{153}\text{Cl}_{20}\text{N}_{30}\text{O}_{31}\text{Na}$  3244; found 3242 for (clp-Clp-Gly) $_{10}$ .

### CD Spectroscopy of (Pro-Clp-Gly) $_7$ , (clp-Pro-Gly) $_7$ , (Pro-Clp-Gly) $_{10}$ , (clp-Pro-Gly) $_{10}$ , (clp-Clp-Gly) $_{10}$ , and (Clp-Pro-Gly) $_{10}$

Peptides were dried under vacuum for at least 24 h before being weighed and dissolved to 0.2 mM in 50 mM acetic acid (pH 3.0). These solutions were incubated at  $\leq 4^\circ\text{C}$  for  $\geq 24$  h before CD spectra were acquired with an Aviv 202SF spectrometer at the University of Wisconsin Biophysics Instrumentation Facility. Spectra were measured with a 1-nm band-pass in cuvettes with a 0.1-cm path-length. The signal was averaged for 3 s during wavelength scans and 5 s during denaturation experiments using a  $0.6^\circ\text{C}$  temperature deadband. During denaturation experiments, CD spectra were acquired at intervals of  $3^\circ\text{C}$ . At each temperature, solutions were equilibrated for 5 min before data acquisition. Values of  $T_m$  were determined by fitting the molar ellipticity at 225 nm for (Pro-Clp-Gly) $_7$  and (clp-Pro-Gly) $_7$  or 226 nm for the other four peptides to a two-state model.<sup>80</sup>  $T_m$  values were determined in triplicate.

### X-Ray Crystallography

The crystals of Ac-Clp-OMe (10) used for X-ray structure determination were obtained by dissolving the colorless oil in a minimum of ethyl acetate and equilibrating with a reservoir of hexanes. Crystals suitable for X-ray crystallography grew slowly over the course of 2 months. The experimental procedure for the structure determination and tables of atomic coordinates, bond lengths, bond angles, and torsion angles are provided in the Supporting Information.

### Sedimentation Equilibrium Experiments on (clp-Pro-Gly) $_{10}$ , (Pro-Clp-Gly) $_{10}$ , and (clp-Clp-Gly) $_{10}$

Sedimentation equilibrium experiments were performed with a Beckman XL-A analytical ultracentrifuge at the University of Wisconsin Biophysics Instrumentation Facility to evaluate the self-assembly of the peptides (clp-Pro-Gly) $_{10}$ , (Pro-Clp-Gly) $_{10}$ , and (clp-Clp-Gly) $_{10}$ . Detailed experimental procedures are provided in the Supporting Information.

This work is dedicated to the memory of Elkan R. Blout (1919–2006). We are grateful to D. Gottlieb, J. Kalia, and D.R. McCaslin for helpful discussion. MDS was supported by a Department of Homeland Security Graduate Fellowship. CD, MALDI–TOF, and sedimentation equilibrium experiments were performed at the University of Wisconsin–Madison Biophysics Instrumentation Facility.

## REFERENCES

1. Myllyharju, J.; Kivirikko, K. I. *Ann Med* 2001, 33, 7–21.
2. Ramachandran, G. N.; Kartha, G. *Nature* 1954, 174, 269–270.
3. Ramachandran, G. N.; Kartha, G. *Nature* 1955, 176, 593–595.
4. Rich, A.; Crick, F. H. C. *Nature* 1955, 176, 915–916.
5. Cowan, P. M.; McGavin, S.; North, A. C. T. *Nature* 1955, 176, 1062–1064.
6. Rich, A.; Crick, F. H. C. *J Mol Biol* 1961, 3, 483–506.
7. Ricard-Blum, S.; Ruggiero, F.; van der Rest, M. *Top Curr Chem* 2005, 247, 35–84.
8. Veit, G.; Kobbe, B.; Keene, D. R.; Paulsson, M.; Koch, M.; Wagener, R. *J Biol Chem* 2006, 281, 3494–3504.
9. Jenkins, C. L.; Raines, R. T. *Nat Prod Rep* 2002, 19, 49–59.
10. Brodsky, B.; Persikov, A. V. *Adv Protein Chem* 2005, 70, 301–339.
11. Bella, J.; Eaton, M.; Brodsky, B.; Berman, H. M. *Science* 1994, 266, 75–81.
12. Fietzek, P. P.; Kühn, K. *Mol Cell Biochem* 1975, 8, 141–157.
13. Engel, J.; Bächinger, H. P. *Top Curr Chem* 2005, 247, 7–33.
14. Myllyharju, J. *Matrix Biol* 2003, 22, 15–24.
15. Friedman, L.; Higgin, J. J.; Moulder, G.; Barstead, R.; Raines, R. T.; Kimble, J. *Proc Natl Acad Sci USA* 2000, 97, 4736–4741.
16. Holster, T.; Pakkanen, O.; Soininen, R.; Sormunen, R.; Nokelainen, M.; Kivirikko, K. I.; Myllyharju, J. *J Biol Chem* 2007, 282, 2512–2519.
17. Berg, R. A.; Prockop, D. J. *Biochem Biophys Res Commun* 1973, 52, 115–120.
18. Burjanadze, T. V. *Biopolymers* 2000, 53, 523–528.
19. Brown, F. R., III; Carver, J. P.; Blout, E. R. *J Mol Biol* 1969, 39, 307–313.
20. Brown, F. R., III; di Corato, A.; Lorenzi, G. P.; Blout, E. R. *J Mol Biol* 1972, 63, 85–99.
21. Doyle, B. B.; Bendit, E. G.; Blout, E. R. *Biopolymers* 1975, 14, 937–957.
22. Sakakibara, S.; Inouye, K.; Shudo, K.; Kishida, Y.; Kobayashi, Y.; Prockop, D. J. *Biochim Biophys Acta* 1973, 303, 198–202.
23. Inouye, K.; Sakakibara, S.; Prockop, D. J. *Biochim Biophys Acta* 1976, 420, 133–141.
24. Inouye, K.; Kobayashi, Y.; Kyogoku, Y.; Kishida, Y.; Sakakibara, S.; Prockop, D. J. *Arch Biochem Biophys* 1982, 219, 198–203.
25. Bretscher, L. E.; Jenkins, C. L.; Taylor, K. M.; DeRider, M. L.; Raines, R. T. *J Am Chem Soc* 2001, 123, 777–778.
26. Hodges, J. A.; Raines, R. T. *J Am Chem Soc* 2003, 125, 9262–9263.
27. Shoulders, M. D.; Hodges, J. A.; Raines, R. T. *J Am Chem Soc* 2006, 128, 8112–8113.
28. Hodges, J. A.; Raines, R. T. *J Am Chem Soc* 2005, 127, 15923–15932.
29. Holmgren, S. K.; Bretscher, L. E.; Taylor, K. M.; Raines, R. T. *Chem Biol* 1999, 6, 63–70.
30. Nishi, Y.; Uchiyama, S.; Doi, M.; Nishiuchi, Y.; Nakazawa, T.; Ohkubo, T.; Kobayashi, Y. *Biochemistry* 2005, 44, 6034–6042.
31. Bann, J. G.; Bächinger, H. P. *J Biol Chem* 2000, 275, 24466–24469.
32. Mizuno, K.; Hayashi, T.; Bächinger, H. P. *J Biol Chem* 2003, 278, 32373–32379.
33. Mizuno, K.; Hayashi, T.; Peyton, D. H.; Bächinger, H. P. *J Biol Chem* 2004, 279, 38072–38078.
34. Kawahara, K.; Nishi, Y.; Nakamura, S.; Uchiyama, S.; Nishiuchi, Y.; Nakazawa, T.; Ohkubo, T.; Kobayashi, Y. *Biochemistry* 2005, 44, 15812–15822.
35. Schumacher, M.; Mizuno, K.; Bächinger, H. P. *J Biol Chem* 2005, 280, 20397–20403.
36. Raines, R. T. *Protein Sci* 2006, 15, 1219–1225.
37. Holmgren, S. K.; Taylor, K. M.; Bretscher, L. E.; Raines, R. T. *Nature* 1998, 392, 666–667.
38. DeRider, M. L.; Wilkens, S. J.; Waddell, M. J.; Bretscher, L. E.; Weinhold, F.; Raines, R. T.; Markley, J. L. *J Am Chem Soc* 2002, 124, 2497–2505.
39. Improta, R.; Benzi, C.; Barone, V. *J Am Chem Soc* 2001, 123, 12568–12577.
40. Giacomazzo, C.; Monaco, H. L.; Artioli, G.; Viterbo, D.; Ferraris, G.; Gilli, G.; Zanotti, G.; Catti, M. *Fundamentals of Crystallography*, 2nd ed.; Oxford University Press: Oxford, UK, 2002.
41. Umashankara, M.; Babu, I. R.; Ganesh, K. N. *Chem Commun* 2003, 2606–2607.
42. Vitagliano, L.; Berisio, R.; Mazzarella, L.; Zagari, A. *Biopolymers* 2001, 58, 459–464.
43. Doi, M.; Nishi, Y.; Uchiyama, S.; Nishiuchi, Y.; Nakazawa, T.; Ohkubo, T.; Kobayashi, Y. *J Am Chem Soc* 2003, 125, 9922–9923.
44. Barth, D.; Milbradt, A. G.; Renner, C.; Moroder, L. *ChemBioChem* 2004, 5, 79–86.
45. Hinderaker, M. P.; Raines, R. T. *Protein Sci* 2003, 12, 1188–1194.
46. Hodges, J. A.; Raines, R. T. *Org Lett* 2006, 8, 4695–4697.
47. Sonntag, L.-S.; Schweizer, S.; Ochsenfeld, C.; Wennemers, H. *J Am Chem Soc* 2006, 128, 14697–14703.
48. Gorske, B. C.; Bastian, B. L.; Geske, G. D.; Blackwell, H. E. *J Am Chem Soc* 2007, 129, 8928–8929.
49. Dunitz, J. D.; Taylor, R. *Chem Eur J* 1997, 3, 89–98.
50. Banerjee, R.; Desiraju, G. R.; Mondal, R.; Howard, J. A. K. *Chem Eur J* 2004, 10, 3373–3383.
51. Pauling, L. *The Nature of the Chemical Bond*; Cornell University Press: Ithaca, NY, 1960.
52. Hansch, C.; Leo, A.; Taft, R. W. *Chem Rev* 1991, 91, 165–195.
53. Alabugin, I. V.; Zeidan, T. A. *J Am Chem Soc* 2002, 124, 3175–3185.
54. Mauger, A. B. *J Nat Prod* 1996, 59, 1205–1211.
55. Yoshioka, H.; Nakatsu, K.; Sato, M.; Tatsuno, T. *Chem Lett* 1973, 1319–1322.
56. Morita, H.; Nagashima, S.; Takeya, K.; Itokawa, H. *Chem Pharm Bull* 1993, 41, 992–993.
57. Morita, H.; Nagashima, S.; Takeya, K.; Itokawa, H. *Tetrahedron* 1994, 50, 11613–11622.
58. Morita, H.; Nagashima, S.; Takeya, K.; Itokawa, H.; Iitaka, Y. *Tetrahedron* 1995, 51, 1121–1132.
59. Tan, N.-H.; Zhou, J. *Chem Rev* 2006, 106, 840–895.
60. Morita, H.; Nagashima, S.; Takeya, K.; Itokawa, H. *Chem Lett* 1994, 2009–2010.
61. Lee, S.; Noda, K.; Aoyagi, H.; Kato, T.; Izumiya, N. *Int J Pept Protein Res* 1986, 27, 44–50.
62. Okumara, Y.; Okamoto, R.; Ishikura, T. *Agric Biol Chem* 1984, 48, 543–544.

63. Mauger, A. B.; Thomas, W. A. *Org Magn Reson* 1981, 17, 186–191.
64. Chiba, J.; Takayama, G.; Takashi, T.; Yokoyama, M.; Nakayama, A.; Baldwin, J. J.; McDonald, E.; Moriarty, K. J.; Sarko, C. R.; Saionz, K. W.; Swanson, R.; Hussain, Z.; Wong, A.; Machinaga, N. *Bioorg Med Chem* 2006, 14, 2725–2746.
65. Appel, R. *Angew Chem Int Ed Engl* 1975, 14, 801–811.
66. Berger, Y.; Dehmlow, H.; Blum-Kaelin, D.; Kitas, E. A.; Löffler, B.-M.; Aebi, J. D.; Juillerat-Jeanneret, L. *J Med Chem* 2005, 48, 483–498.
67. Panasik, N., Jr.; Eberhardt, E. S.; Edison, A. S.; Powell, D. R.; Raines, R. T. *Int J Pept Protein Res* 1994, 44, 262–269.
68. Eberhardt, E. S.; Panasik, N., Jr.; Raines, R. T. *J Am Chem Soc* 1996, 118, 12261–12266.
69. Jenkins, C. L.; McCloskey, A. I.; Guzei, I. A.; Eberhardt, E. S.; Raines, R. T. *Biopolymers: Pept Sci* 2005, 80, 1–8.
70. Nudelman, A.; Bechor, Y.; Falb, E.; Fischer, B.; Wexler, B. A.; Nudelman, A. *Synth Commun* 1998, 28, 471–474.
71. Coste, J.; Le-Nguyen, D.; Castro, B. *Tetrahedron Lett* 1990, 31, 205–208.
72. Coste, J.; Frérot, E.; Jouin, P.; Castro, B. *Tetrahedron Lett* 1991, 32, 1967–1970.
73. Doi, M.; Nishi, Y.; Uchiyama, S.; Nishiuchi, Y.; Nishio, H.; Nakazawa, T.; Ohkubo, T.; Kobayashi, Y. *J Pept Sci* 2005, 11, 609–616.
74. Horng, J.-C.; Raines, R. T. *Protein Sci* 2006, 15, 74–83.
75. Ruzza, P.; Siligardi, G.; Donella-Deana, A.; Calderan, A.; Hussain, R.; Rubini, C.; Cesaro, L.; Osler, A.; Guiotto, A.; Pinna, L. A.; Borin, G. *J Pept Sci* 2006, 12, 462–471.
76. Myllyharju, J. *Top Curr Chem* 2005, 247, 115–147.
77. Blasiak, L. C.; Vaillancourt, F. H.; Walsh, C. T.; Drennan, C. L. *Nature* 2006, 440, 368–371.
78. Vaillancourt, F. H.; Yeh, E.; Vosburg, D. A.; Garneau-Tsodikova, S.; Walsh, C. T. *Chem Rev* 2006, 106, 3364–3378.
79. Jenkins, C. L.; Vasbinder, M. M.; Miller, S. J.; Raines, R. T. *Org Lett* 2005, 7, 2619–2622.
80. Becktel, W. J.; Schellman, J. A. *Biopolymers* 1987, 26, 1859–1877.

*Reviewing Editor: Lila Gierasch*