

Effect of Bovine Seminal Ribonuclease and Bovine Pancreatic Ribonuclease A on Bovine Oocyte Maturation

TOMÁŠ SLAVÍK,^{1*} JOSEF MATOUŠEK,¹ JOSEF FULKA,¹ AND RONALD T. RAINES²

¹*Institute of Animal Physiology and Genetics, Academy of Sciences of the Czech Republic, Czech Republic*

²*Department of Biochemistry, University of Wisconsin, Madison, Wisconsin 53706-1569*

ABSTRACT Bovine seminal ribonuclease (BS-RNase) contains the M×M (noncovalent dimer) and M=M (free monomer) in constant ratio. The aim of this work was to evaluate the effect of BS-RNase, its monomer and dimer forms, and also various mutants of this enzyme on meiotic completion in cattle oocytes. It was found that BS-RNase has irreversible effects on the meiotic maturation of bovine oocytes *in vitro*, particularly on the completion of meiosis. The effect of BS-RNase is dose-dependent. In medium supplemented with 1 µg/ml, the results were comparable with those of the control (70% MII oocytes after 24 hr of culture). Whereas 5 µg/ml reduced the number of MII oocytes to 50%, 10 and 25 µg/ml arrested this process completely. The M×M form and RNase A at 5 µg/ml inhibited the maturation rate by 71 and 48%, respectively, but a less significant effect was observed for the M=M form, or the carboxymethylated monomers MCM31 and MCM32 (21%, 16%, and 42% MII oocytes, respectively, in comparison with control). These data demonstrate that bovine ribonucleases can have variable detrimental effects on the maturation of bovine oocyte. *J. Exp. Zool.* 287:394–399, 2000. © 2000 Wiley-Liss, Inc.

Bovids have high levels of two homologous secretory ribonucleases (EC 3.1.27.5): bovine seminal ribonuclease (BS-RNase) and bovine pancreatic ribonuclease A (RNase A). These enzymes can have significant effects on animal physiology. For example, the aspermatogenic activity of BS-RNase has been demonstrated in males from several species (Matoušek et al., '73; Dostál and Matoušek, '73; Leone et al., '73; Matoušek et al., '78; Hlinak et al., '81; Matoušek, '94). Preliminary investigations of the embryotoxic effect of BS-RNase have been carried out in female mice, guinea pigs, rabbits, and pigs (Matoušek and Grozdanovič '73; Matoušek et al., '73a,b). Here we continue this analysis and report on the effects of wild-type and mutant forms of BS-RNase and RNase A on bovine oocyte maturation.

BS-RNase has been prepared from a synthetic gene in *Escherichia coli* (Kim and Raines, '93). This preparation has the same antiembryonic activity as that of BS-RNase isolated from seminal plasma (Kim et al., '95a). In addition, mutant forms of the recombinant enzyme are embryotoxic (Kim et al., '95a). The cytotoxicity of BS-RNase is related to its quaternary structure. BS-RNase is isolated

as a dimer that is cross-linked by two disulfide bonds. Monomeric BS-RNase and bovine pancreatic ribonuclease A (RNase A), which is a monomer, are not cytotoxic (Vescia et al., '80; Tamburini et al., '90; Kim et al., '95a). In contrast, artificially dimerized RNase A is cytotoxic but to a lesser extent than is BS-RNase (Bartholeyns and Baudhuin, '76; Bartholeyns and Zenebergh, '79; Vescia et al., '80; Di Donato et al., '94).

At equilibrium, dimeric BS-RNase is a mixture of two distinct quaternary forms, M=M and M×M (Piccoli et al., '92). The conversion of M=M to M×M entails the exchange of N-terminal helices between subunits, as occurs during the artificial dimerization of RNase A. We and others have demonstrated that it is the M×M form that is respon-

Grant sponsor: Grant Agency of the Czech Republic; Grant numbers: 523/96/0710 and 524/99/0844; Grant sponsor: National Agency for Agricultural Research NAZV CR; Grant number: EP0960006203; Grant sponsor: European Commission; Grant number: PL 966052; Grant sponsor: National Institutes of Health; Grant number: GM44783.

*Correspondence to: T. Slavík, Institute of Animal Physiology and Genetics, Academy of Sciences of the Czech Republic, 277 21 Liběchov, Czech Republic. E-mail: slavik@iapg.cas.cz

Received 5 August 1998; Accepted 27 March 2000

sible for the cytotoxicity of BS-RNase (Cafaro et al., '95; Kim et al., '95a). We have also demonstrated that the ribonucleolytic activity of BS-RNase is necessary for its cytotoxic activity (Kim et al., '95a).

Thirty years ago, Roth ('67) discovered that mammalian cells contain ribonuclease inhibitor (RI), a protein that binds tightly to monomeric but not dimeric BS-RNase (Murthy and Sirdeshmukh, '92; Kim et al., '95b; Murthy et al., '96). RI appears to act as a sentry that protects cellular RNA from invading secretory ribonucleases (Blackburn and Moore '82; Hofsteenge, '94). A key to the cytotoxicity of BS-RNase is likely to be its ability to evade cellular RI (Cafaro et al., '95; Di Donato et al., '95; Kim et al., '95b,c).

Monomers of RNase A, like those of BS-RNase, are bound tightly by RI and are not cytotoxic. Onconase, which is a monomeric ribonuclease isolated from the eggs of the Northern leopard frog (*Rana pipiens*) resists RI and is cytotoxic (Ardelt et al., '91; Wu et al., '93; Boix et al., '96). X-ray diffraction analysis revealed that extensive contacts between RNase A and RI occur in the region near Asp31 of RI and Lys31 of RNase A (Kobe and Deisenhofer, '95). This contact appears to contribute to the stability of the RI-RNase A complex. Here we tested the effect of MCM31 and MCM32, which are monomers with a carboxymethyl group ($-\text{CH}_2\text{CO}_2-$) on Cys31 or Cys32 (Matoušek et al., '96), and other ribonucleases on bovine oocyte maturation *in vitro*.

MATERIALS AND METHODS

Preparation of ribonucleases

Wild-type BS-RNase was isolated from seminal vesicle fluid by precipitation with acetic acid and ammonium sulfate and subjected to chromatography on columns of CM Sephadex C-50 and Sephadex G-100 (Dostál and Matoušek, '73). The purity of BS-RNase was assessed by peak homogeneity during chromatography; by disc electrophoresis in an acrylamide gel; by starch-gel electrophoresis; by immunoelectrophoresis in an agar gel; and by ultracentrifugation (Dostál and Matoušek, '73). Recombinant BS-RNase was produced in *E. coli* as described (Kim and Raines, '93).

Two distinct quaternary forms of BS-RNase dimer, designated as M×M and M=M (Piccoli et al., '92), were purified from bull seminal plasma as described (Tamburrini et al., '86). Briefly, the inter-subunit disulfide bonds of the purified en-

zyme were reduced with a 10-fold molar excess of reduced dithiothreitol, and the resulting protein was subjected to gel filtration chromatography to separate monomer from the noncovalent dimer. The M=M form was prepared by air oxidation of monomer and purified by gel filtration chromatography. Similarly, the M×M form was prepared by air oxidation of noncovalent dimer and purified by gel filtration chromatography. This method of purification yielded M=M and M×M that were >90% free of the other form, as judged by selective reduction followed by gel filtration chromatography (Kim et al., '95a).

Oligonucleotide-mediated site-directed mutagenesis was used to change the TGT codon of Cys31 or Cys32 to the TCT codon of serine as described (Kim et al., '95a). The resulting plasmids, pLSR31 and pLSR32, were used to produce C31S BS-RNase and C32S BS-RNase, respectively. MCM31 and MCM32 BS-RNase monomers were prepared from C32S and C31S BS-RNase as described (Matoušek et al., '96).

Assay of oocyte maturation in vitro

Cattle oocytes were obtained from ovaries of normally cycling animals after slaughter. The ovaries were transported at about 20°C to the laboratory within 2 hr, and once there they were rinsed briefly with 95% (v/v) ethanol and repeatedly with saline solution containing antibiotics (streptomycin and penicillin). Ovaries were cut and scarified under phosphate-buffered saline containing heparin (100 IU/ml) and isolated under a stereomicroscope. Oocytes were evaluated, and only those with a compact cumulus were used for culture under an atmosphere of N₂ (85% v/v), O₂ (10% v/v), and CO₂ (5% v/v) at 38.5°C for 24–26 hr (Kubelka et al., '88). The composition of culture medium was the same as described by Pavlok et al. ('88). This medium contained 80 ml of TC 199 (Sevac, Prague), 39 ml of 7.5% (w/v) NaHCO₃ (Sevac, Prague), 1,500 mg of HEPES (acid form; Serva, Heidelberg), 600 mg of lactic acid (calcium salt; Serva, Heidelberg), 200 mg of pyruvic acid (sodium salt; Serva, Heidelberg), 25 IU/ml penicillin (potassium salt), and 25 IU of streptomycin sulfate, and was made up to 1.00 l with Nanopure-filtered water. Before it was used for culture, the medium was supplemented with bovine serum (BOS; 15% v/v), FSH (0.2 IU/ml; Calbiochem, La Jolla, CA) and with one of selected ribonucleases in the concentration 0, 1, 5, 10, or 25 µg/ml.

To test the reversibility of a ribonuclease effect, the ribonuclease-containing medium was replaced

after 24 hr of culture with ribonuclease-free medium. The oocytes were then cultured for an additional 24 hr (Kubelka et al., '88).

At the beginning and at the end of a culture period, some of the oocytes were removed for nuclear evaluation. After fixation in acetalcohol overnight, the oocytes were stained with aceto-orcein (2% w/v) and examined by phase-contrast microscopy. Between 14 and 52 oocytes were examined for each type of maturation medium.

The proportion of MII oocytes with extruded polar bodies was used as a unique criterion of maturation. The suitability of this procedure had been demonstrated previously in experiments in which oocytes matured under the identical culture protocol described above resulted in healthy progeny after *in vitro* fertilization and transfer to foster mothers (Pavlok et al., '88; Slavík and Fulka, '91; Slavík et al., '92).

Fisher's exact test was used for statistical evaluation.

RESULTS

Oocyte maturation in vitro

Maturation rates of oocytes in the each experimental group were compared with meiotic progression to MII in controls. The effect of wild-type BS-RNase was dose dependent (Fig. 1). Oocyte maturation was not affected significantly by 1 μ g of BS-RNase per ml. Rather, maturation was comparable with that of control oocytes (70% MII oocytes after 24 hr of culture), and no degeneration

was observed under a phase-contrast microscope. In contrast, 5 μ g/ml reduced the maturation rate by approximately 1/3 in comparison with controls (70% vs. 43%); 10 and 25 μ g/ml arrested this process completely. Moreover, at higher concentrations of BS-RNase, remarkable degenerative changes in the cytoplasm and an atypical configuration of chromatin were observed. In these two groups about half of oocytes had been ruptured during routine fixation, most likely due to disintegration of cytoskeletal structures, and thus they were included in the category of degenerated oocytes. (Fig. 1).

The effect of BS-RNase on oocyte maturation was not reversible. No resumption of the maturation process was observed after replacing the medium containing 5 μ g of BS-RNase per ml with ribonuclease-free medium (Fig. 2).

Marked differences were found among the mutant ribonuclease tested. In medium supplemented with BS-RNase, its M \times M form, and RNase A at 5 μ g/ml, the maturation rate was significantly (** $P > 0.01$, * $P > 0.05$) inhibited by 45% (*), 48% (*), and 71% (**), respectively, in comparison with a control group (100%). On the other hand, a statistically non-significant effect was observed for the M=M form, MCM31, and MCM32 (21%, 16%, and 42%, respectively) in comparison with a control group (Fig. 3).

From these results it may be concluded that ribonucleases and their mutants affected resumption of meiosis of bovine oocytes and particularly the completion of this process.




Fig. 1. Effect of added BS-RNase concentration on maturation of bovine oocytes *in vitro*. Data are the mean

values for 24 hr incubation of 14–34 oocytes per group (2 replications).




Fig. 2. Reversibility of inhibitory effect of BS-RNase on maturation of bovine oocytes in vitro. The concentration of BS-RNase was 5 µg per ml for 24 hr (inhibition) and 0 µg per

ml for additional 24 hr (reversibility). The control cultures contained no added ribonuclease. Data are the mean values for 23–30 oocytes per group (3 replications).

DISCUSSION

In this work, we evaluated the effect of BS-RNase and variants of this enzyme on final period of bovine oocyte formation. The variants were chosen with the particular aim of correlating their quaternary structure with cytotoxic activities during the final part of meiosis.

Bovine seminal ribonuclease at a concentration of 5 µg/ml reduces the rate of bovine oocyte

maturation to the MII stage by approximately 58%. Ten and 25 µg/ml stop maturation completely (Fig. 1). Further, the higher concentrations of BS-RNase increases markedly the number of oocytes with degenerative cytoplasmic changes and chromatin disorders.

Mammalian oocyte together with surrounding cumulus cells represents specific experimental model. During growth and meiotic maturation,




Fig. 3. Effect of ribonucleases on the maturation of bovine oocytes in vitro. The concentration of added ribonuclease was 5 µg per ml. The control cultures contained no added ribonuclease. Data are the mean values for 24 hr incubations

of 31–52 oocytes per group (3 replications). Statistically significant differences in comparison with control: ***P* > 0.01, **P* > 0.05.

oocyte and cumulus cells form functional complexes that are characteristic with mutual interaction between these two cell categories (Eppig, '91). While RNA synthesis in fully grown oocytes, which were used in our experiments, is very low (Crozet et al., '86), cumulus cells behave as normal mitotic cells. Under routinely used culture protocol and also in vivo, they supply the oocyte with nutritional substances and with not yet defined molecular signals, which are responsible for the regulation of meiotic maturation. Therefore, it cannot be excluded that the inhibition of meiosis and degenerative changes in maturing oocytes were caused indirectly as a toxic effect of BS-RNase or its variants upon cumulus cells which were not able to provide oocytes with adequate array of molecules.

In previous work, we showed that the M×M form of BS-RNase has significantly higher cytotoxic activity than did the M=M form (Kim et al., '95b). We therefore proposed that the difference in the cytotoxicity of the two forms derives from their different fates inside the cell. More specifically, the basis of the different activities of M×M and M=M is the ability of M×M but not M=M to remain dimeric in a reducing environment, such as the mammalian cytosol.

In contrast to its relative low cytotoxicity, the M=M form of BS-RNase has significant anti-meiotic activity in vitro, similar to that of the M×M form (Fig. 3). The antimetabolic effects as well as antitumor and immunosuppressive activities of the M=M form (Kim et al., '95b) were demonstrated in vitro, where identical cell-cell interactions do not exist. This result contrasts somewhat with a modest aspermatogenic effect reported recently (Kim et al., '95b). The aspermatogenic activity was, however, assessed in vivo, where more complex interactions might influence degradation or inhibition of ribonucleases.

Until this decade, only dimeric ribonucleases were known to have significant cytotoxic activity (Bartholeyns and Baudhuin, '76; Bartholeyns and Zeneberg, '77; Di Donato et al., '94, '95). The demonstrated cytotoxicity of onconase (Ardelt et al., '91; Wu et al., '93, '95) and angiogenin (Matoušek et al., '95), two monomeric ribonucleases, reveals that a dimeric form is not essential for a ribonuclease to be cytotoxic. The aspermatogenic activity of MCM31 and MCM32 (Matoušek et al., '96), which have no effect on oocyte maturation in assays in which other monomeric ribonucleases are active, casts further doubt on the relationship between a dimeric form and cytotoxicity. The key to

ribonuclease cytotoxicity appears to be the evasion of cellular ribonuclease inhibitor RI (Cafaro et al., '95; Di Donato et al., '95; Kim et al., '95a,b). Different tissues contain different levels of RI (Lee and Vallee, '93). Thus, a possible explanation for the toxicity of MCM31 and MCM32 for spermatogenic layers is that these cells do not contain enough RI to inactivate invading MCM31 and MCM32. From this perspective, the bovine ova have enough RI to block MCM31 and MCM32. Nevertheless, no explanation is apparent for the sensitivity of these cells to RNase A, which is also a monomer. Perhaps differential adsorption of ribonucleases also plays a role in the cell-specificity of cytotoxicity. Testing this hypothesis will be the basis of the future work.

ACKNOWLEDGMENTS

This research was supported by grants 523/96/0710 (to J.M.) and 524/99/0844 (to T.S.) from the Grant Agency of the Czech Republic, grant EP096-0006203 from National Agency for Agricultural Research NAZV CR (to T.S.), grant PL 966052 from the European Commission (to J.M.), and grant GM44783 from the National Institutes of Health (to R.T.R.). The authors thank M. Hokešová for technical collaboration.

LITERATURE CITED

- Ardelt W, Mikulski SM, Shogen K. 1991. Amino acid sequence of an antitumor protein from *Rana pipiens* oocytes and early embryos. *J Biol Chem* 266:245-251.
- Bartholeyns J, Baudhuin P. 1976. Inhibition of tumor cell proliferation by dimerized ribonuclease. *Proc Natl Acad Sci USA* 73:573-576.
- Bartholeyns J, Zenebergh A. 1979. In vitro and in vivo anti-tumor effect of dimerized ribonuclease A. *Eur J Cancer* 15:85-91.
- Blackburn P, Moore S. 1982. Pancreatic ribonuclease. In: Boyer PD, editor. *The enzymes*, vol 15. New York: Academic Press. p 317-433.
- Boix E, Wu Y, Vasandani VM, Saxena SK, Ardel W, Ladner J, Youle RJ. 1996. Role of the N terminus in RNase A homologues: differences in catalytic activity, ribonuclease inhibitor interaction and cytotoxicity. *J Mol Biol* 257:992-1007.
- Cafaro V, De Lorenzo C, Piccoli R, Bracale A, Mastronicola MR, Di Donato A, D'Alessio G. 1995. The antitumor action of seminal ribonuclease and its quaternary conformations. *FEBS Lett* 359:31-34.
- Crozet N, Motlik J, Kaňka J, Fulka J. 1986. Nucleolar fine structure and RNA synthesis in bovine oocytes from antral follicles. *Gamete Res* 14:65-73.
- D'Alessio G, Malorni MC, Parente A. 1975. Dissociation of bovine seminal ribonuclease into catalytically active monomers by selective reduction and alkylation of the inter-subunit disulfide bridges. *Biochemistry* 14:1116-1122.
- de Nigris M, Russo N, Piccoli R, D'Alessio G, Di Donato A. 1993. Expression of bovine seminal ribonuclease in *Escherichia coli*. *Biochem Biophys Res Commun* 193:155-160.

- Di Donato A, Cafaro V, D'Alessio G. 1994. Ribonuclease A can be transformed into a dimeric ribonuclease with anti-tumor activity. *J Biol Chem* 269:17394–17396.
- Di Donato A, Cafaro V, Romeo I, D'Alessio G. 1995. Hints on the evolutionary design of a dimeric RNase with special bioactions. *Protein Sci* 4:1470–1477.
- Dostál J, Matoušek J. 1973. Isolation and some chemical properties of aspermatogenic substance from bull seminal vesicle fluid. *J Reprod Fertil* 33:263–274.
- Guillemette JG, Matsushima-Hibiya Y, Atkinson T, Smith M. 1991. Expression in *Escherichia coli* of a synthetic gene coding for horse heart myoglobin. *Protein Eng* 4:585–592.
- Eppig JJ. 1991. Intercommunication between mammalian oocytes and companion somatic cells. *BioEssays* 11:569–574.
- Hlinak Z, Matoušek J, Madlafousek J. 1981. The effect of bull seminal ribonuclease on reproductive organs and sexual behaviour in male rats. *Physiol Bohemoslov* 30:539–542.
- Hofsteenge J. 1994. "Holy" proteins I: ribonuclease inhibitor. *Curr Opin Struct Biol* 4:807–809.
- Kim J-S, Raines RT. 1993. Bovine seminal ribonuclease produced from a synthetic gene. *J Biol Chem* 268:17392–17396.
- Kim J-S, Souček J, Matoušek J, Raines RT. 1995a. Catalytic activity of bovine seminal ribonuclease is essential for its immunosuppressive and other biological activities. *Biochem J* 308:547–550.
- Kim J-S, Souček J, Matoušek J, Raines RT. 1995b. Structural basis for the biological activities of bovine seminal ribonuclease. *J Biol Chem* 270:10525–10530.
- Kim J-S, Souček J, Matoušek J, Raines RT. 1995c. Mechanism of ribonuclease cytotoxicity. *J Biol Chem* 270:31097–31102.
- Kobe B, Deisenhofer J. 1995. A structural basis of the interactions between leucine-rich repeats and protein ligands. *Nature* 374:183–186.
- Kubelka M, Motlík J, Fulka J Jr, Procházka R, Rimkevičová Z, Fulka J. 1988. Time sequence of germinal vesicle breakdown in pig oocytes after cycloheximide and *p*-aminobenzamide block. *Gamete Res* 19:423–431.
- Leone E, Greco L, Rastogi RK, Iela L. 1973. Antispermatic properties of bull seminal ribonuclease. *J Reprod Fertil* 34:197–200.
- Matoušek J. 1973. Antiembryonic effect of bull seminal vesicle fluid in mice. *J Reprod Fertil* 32:175–184.
- Matoušek J. 1994. Aspermatogenic effect of the bull seminal ribonuclease (BS RNase) in the presence of anti-BS RNase antibodies in mice. *Anim Genet* 25(Suppl 1):45–50.
- Matoušek J, Grozdanović J. 1973. Specific effect of bull seminal ribonuclease (AS RNase) on cell systems in mice. *Comp Biochem Physiol* 46A:241–248.
- Matoušek J, Pavlok A, Dostál J, Grozdanović J. 1973a. Some biological properties of bull seminal vesicle aspermatogenic substance and its effect on mice. *J Reprod Fertil* 34:9–22.
- Matoušek J, Fulka J, Pavlok A. 1973b. Effect of ribonuclease fractions isolated from bull seminal vesicle fluid on embryonic mortality in guinea pigs, rabbits and pigs. *Int J Fertil* 18:13–16.
- Matoušek J, Souček J, Říha J, Zankel TR, Benner SA. 1995. Immunosuppressive activity of angiogenin in comparison with bovine seminal ribonuclease and pancreatic ribonuclease. *Comp Biochem Physiol* 112B:235–241.
- Matoušek J, Kim J-S, Souček J, Říha J, Ribo M, Leland PA, Raines RT. 1997. Ribonucleases endowed with specific toxicity for spermatogenic layers. *Comp Biochem Physiol B* 118B:887–888.
- Matoušek J, Veselský L, Baše J, Dostál J. 1978. The effect of bull seminal ribonuclease on spermatogenesis in the rabbit. *Anim Reprod Sci* 1:49–59.
- Murthy BS, De Lorenzo C, Piccoli R, D'Alessio G, Sirdeshmukh R. 1996. Effects of protein RNase inhibitor and substrate on the quaternary structures of bovine seminal RNase. *Biochemistry* 35:3880–3885.
- Murthy BS, Sirdeshmukh R. 1992. Sensitivity of monomeric and dimeric forms of bovine seminal ribonuclease to human placental ribonuclease inhibitor. *Biochem J* 281:343–348.
- Pavlok A, Torner H, Motlík J, Fulka J, Kauffold P, Duschinski U. 1988. Fertilization of bovine oocytes in vitro: effect of different sources of gametes on fertilization rate and frequency of fertilization anomalies. *Anim Reprod Sci* 16:207–213.
- Piccoli R, Tamburrini M, Piccialli G, Di Donato A, Parente A, D'Alessio G. 1992. The dual-mode quaternary structure of seminal RNase. *Proc Natl Acad Sci USA* 89:1870–1874.
- Roth JS. 1967. Some observations on the assay and properties of ribonuclease in normal and tumor tissues. *Methods Cancer Res* 3:153–242.
- Slavík T, Fulka J. 1991. Pregnancies after transfer of sheep embryos produced from oocytes matured and fertilized in vitro. *Folia Biol (Praha)* 37:95–99.
- Slavík T, Fulka J, Goll I. 1992. Pregnancy rate after the transfer of sheep embryos originated from randomly chosen oocytes matured and fertilized in vitro. *Theriogenology* 38:749–756.
- Tamburrini M, Piccoli R, De Prisco R, Di Donato A, D'Alessio G. 1986. Fast and high-yielding procedures for the isolation of bovine seminal RNase. *Ital J Biochem* 35:22–32.
- Tamburini M, Scala G, Verde C, Ruocco MR, Parente A, Venuta S, D'Alessio G. 1990. Immunosuppressive activity of bovine seminal RNase on T-cell proliferation. *Eur J Biochem* 190:145–148.
- Vescia S, Tramontano A, Augusti-Tocco G, D'Alessio G. 1980. In vitro studies on selective inhibition of tumor cell growth by seminal ribonuclease. *Cancer Res* 40:3740–3747.
- Wu Y, Mikulski SM, Ardel W, Rybak SM, Youle RJ. 1993. A cytotoxic ribonuclease. *J Biol Chem* 268:10686–10693.
- Wu Y, Saxena SK, Ardel W, Gadina M, Mikulski SM, De Lorenzo V, D'Alessio G, Youle RJ. 1995. A study of the intracellular routing of cytotoxic ribonucleases. *J Biol Chem* 270:17476–17481.